arXiv:1007.1932v4 [math.OA] 23 Nov 2011

NON-COMMUTATIVE FUNCTIONS AND NON-COMMUTATIVE FREE LEVY-HINCIN FORMULA

MIHAI POPA AND VICTOR VINNIKOV

Abstract. The paper is discussing infinite divisibility in the setting of operator-valued boolean, free and, more general, c-free independences. Particularly, using Hilbert bimodules and non-commutative functions techniques, we obtain analogues of the Levy-Hincin integral representation for infinitely divisible real measures.

1. Introduction and notations

The paper presents some results concerning infinite divisibility in the framework of operator-valued non-commutative probability.

In probability theory - classic and non-commutative - limit theorems play a central role. The "most general" limit theorems involve so-called infinitesimal arrays, and the limit distributions are usually identified with *infinitely divisible* distributions. There is a consistent literature about infinitely divisible measures in classical probability (see [9]), dating back to Kolmogorov ([10]), P. Levy ([14]) etc. Similar results have been found for non-commutative independences, such as [3] and [4] for free independence, [23] for boolean probability and [12], [31] for conditionally free probability. In the operator-valued case, when states are replaced by positive conditional expectations or, more general, by completely positive maps between C*- or operator algebras, very little was known about operator-valued infinite divisibility; the only exception we know of was Speicher's work [24]. One of the obstructions is that while in the scalar case important results characterizing infinite divisibility are coming from Nevalinna-Pick representation properties of the functions that linearize additive convolutions (such as the log of the Fourier transform in the classic case or the Voiculescu's R- and ϕ-transforms in the free case), such analytic tools are not yet available in the operator-valued case. The new topic of non-commutative ([11]) or completely matricial ([29]) functions may possibly fill this gap. In particular, the results from Section 5 of the present paper indicate that results similar to the Nevalinna-Pick representation hold also for non-commutative functions.

The paper is organized in five sections. First section presents the introduction and some notations. Sections 2-4 are aimed towards results characterizing infinite divisibility in operator-valued non-commutative probability using combinatorial and operator-algebras methods and constructions. More precisely, in Section 2 we use a non-commutative version of the "Boolean Fock space" construction from [1] to prove that, as in the scalar case, boolean infinite divisibility is trivial in the operator-valued case; particularly, any completely positive map between two C*-algebras is boolean infinitely divisible. Section 3 is describing infinite divisibility with respect to free independence over a positive conditional expectation in terms of maps satisfying a condition of complete positivity. Section 4 is utilizing the techniques from the Boolean case (Section 2) to extend the results of Section 3 from positive conditional expectations to completely positive maps. In particular, we present a construction of the non-commutative version of the conditionally free R-transform of Bozejko, Leinert and Speicher in terms of creation, annihilation and preservation operators on a certain inner-product bimodules. In the scalar-valued case this construction gives a new, combinatorial proof of the main result from [12] (see also [31]) characterizing conditionally free infinite divisibility. In Section 5 we use the tools from the theory of non-commutative functions (see [11], [29]) to define the non-commutative R- (also constructed in [29]), ${}^cR$- and B-transforms. Reformulated in terms of these transforms, the results from Sections 2, 3 and 4 are very similar to the free and conditionally free versions of the Levy-Hincin formula from [3], respectively [12] and [31]. The present material is using the notions detailed in [11] (work in progress), but it is self-contained in this regard, the needed material on non-commutative functions is briefly discussed in Section 5.2.

We will introduce now several notations. Throughout the paper $\mathcal{B}$ will be a unital C*-algebra. We will denote by $\mathcal{B}\langle\mathcal{X}\rangle$ the $*$-algebra freely generated by $\mathcal{B}$ and the selfadjoint symbol $\mathcal{X}$. Unless otherwise explicitly stated, we do not suppose that $\mathcal{B}$ commutes with $\mathcal{X}$. We will also use the notations $\mathcal{B}\langle X\rangle_0$ for the $*$-subalgebra of $\mathcal{B}\langle\mathcal{X}\rangle$ of all polynomials without a free term, and the notation $\mathcal{B}\langle\mathcal{X}_1,\mathcal{X}_2,\dots\rangle$ for the $*$-algebra freely generated by $\mathcal{B}$ and the non-commutating self-adjoint symbols $\mathcal{X}_1,\mathcal{X}_2,\dots$.

In several instances we will identify $\mathcal{T}(\mathcal{B})$, the tensor algebra over $\mathcal{B}$, to the subalgebra of $\mathcal{B}\langle\mathcal{X}\rangle$ spanned by $\{\mathcal{X}b_1\mathcal{X}b_2\cdots\mathcal{X}b_n\colon n\in\mathbb{N}, b_1,\dots,b_n\in\mathcal{B}\}$ via

$$b_1\otimes b_2\otimes\cdots\otimes b_n\mapsto\mathcal{X}b_1\mathcal{X}b_2\cdots\mathcal{X}b_n.$$

The set of all positive conditional expectations from $\mathcal{B}\langle\mathcal{X}\rangle$ to $\mathcal{B}$ will be denoted by $\Sigma_{\mathcal{B}}$.

For $\mathcal{B}\subseteq\mathcal{D}$ a unital inclusion of C*-algebras, we denote by $\Sigma_{\mathcal{B}:\mathcal{D}}$ the set of all unital, positive $\mathcal{B}$-bimodule maps $\mu:\mathcal{B}\langle\mathcal{X}\rangle\longrightarrow\mathcal{D}$ with the property that for all positive integers n and all $\{f_i(\mathcal{X})\}_{i=1}^n\subset\mathcal{B}\langle\mathcal{X}\rangle$ we have that:

$$\left[\mu(f_j(\mathcal{X})^*f_i(\mathcal{X}))\right]_{i,j=1}^n\geq 0 \text{ in } M_n(\mathcal{D}). \tag{1}$$

Remark that $\Sigma_{\mathcal{B}}=\Sigma_{\mathcal{B}:\mathcal{B}}$, as an easy consequence of Exercise 3.18 from [16].

We will denote by $\Sigma_{\mathcal{B}}^0$, respectively $\Sigma_{\mathcal{B}:\mathcal{D}}^0$, the set of all $\mu\in\Sigma_{\mathcal{B}}$ (respectively $\mu\in\Sigma_{\mathcal{B}:\mathcal{D}}$ whose moments do not grow faster than exponentially, that is there exists some $M>0$ such that for all $b_1,\dots,b_n\in\mathcal{B}$, we have

$$||\mu(\mathcal{X}b_1\mathcal{X}b_2\cdots\mathcal{X}b_n\mathcal{X})||<M^{n+1}||b_1||\cdots||b_n||. \tag{2}$$

We will also use the following definition (see [13]):

Definition 1.1. *Let A be a C*-algebra. A* semi-inner-product A-module *is a linear space E which is a right A-module together with a map $(x,y)\mapsto\langle x,y\rangle: E\times E\longrightarrow A$ such that ($x,y,z\in E, a\in A,\alpha,\beta\in\mathbb{C}$):*

(i) $\langle\alpha x+\beta y,z\rangle=\alpha\langle x,y\rangle+\beta\langle y,z\rangle$
(ii) $\langle xa,y\rangle=\langle x,y\rangle a$
(iii) $\langle x,y\rangle=\langle y,x\rangle^*$
(iv) $\langle x,x\rangle\geq 0$

The set of all adjointable maps $T:E\longrightarrow E$ will be denoted by $\mathcal{L}(E)$. Since $\langle\cdot,\cdot\rangle$ is not strictly positive, the adjoint of a map from $\mathcal{L}(E)$ is in general not unique. If

E is a *Hilbert A-module*, that is the inequality at (iv) is strict and E is complete with respect to the norm $\xi \mapsto ||\langle \xi, \xi\rangle||^{\frac{1}{2}}$, then the adjoint is unique and the bounded elements of $\mathcal{L}(E)$ form a C*-algebra.

If $\mathcal{B} \subseteq \mathcal{A}$, $\mathcal{B} \subseteq \mathcal{D}$ are unital inclusions of C*-algebras, $\phi : \mathcal{A} \longrightarrow \mathcal{D}$ is a unital positive $\mathcal{B}$-bimodule map and a is a selfadjoint element of $\mathcal{A}$, we will denote by $\mathcal{B}\langle a\rangle$ the $*$-algebra generated in $\mathcal{A}$ by $\mathcal{B}$ and a and by μ_a, "the $\mathcal{D}$-distribution" of a, that is the positive $\mathcal{B}$-bimodule map $\phi_a : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}$ defined by $\phi_a = \phi \circ \tau_a$ where $\tau_a : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{A}$ is the unique homomorphism such that $\tau_a(\mathcal{X}) = a$ and $\tau_a(b) = b$ for all $b \in \mathcal{B}$. The set of elements from $\Sigma_{\mathcal{B}:\mathcal{D}}$ that can be realized in such way is $\Sigma^0_{\mathcal{B}:\mathcal{D}}$, more precisely we have the following property:

Proposition 1.2. *Let $\mu \in \Sigma_{\mathcal{B}:\mathcal{D}}$. Then $\mu \in \Sigma^0_{\mathcal{B}:\mathcal{D}}$ if and only if there exist a C*-algebra A containing $\mathcal{B}$ as a C*-subalgebra, a completely positive $\mathcal{B}$-bimodule map $\phi : A \longrightarrow \mathcal{D}$ and a self-adjoint element $a \in A$ such that $\mu = \phi_a$.*

Moreover, the condition (2) *is equivalent to the existence of $M > 0$ such that for all $b_1, \ldots, b_n \in M_m(\mathcal{B})$ we have*

$$||(Id_m \otimes \mu)(\mathcal{X} \cdot b_1\mathcal{X} \cdot b_2 \cdots \mathcal{X} \cdot b_n\mathcal{X})|| < M^{n+1}||b_1|| \cdots ||b_n||. \tag{3}$$

where $\mathcal{X}$ acts on $M_m(\mathbb{C}) \otimes \mathcal{B}\langle\mathcal{X}\rangle$ by multiplication on each entry (that is we identify $\mathcal{X}$ to $Id_m \otimes \mathcal{X}$).

Proof. If $\mu = \phi_a$ as above, then the result is trivial.

Suppose now that $\mu \in \Sigma^0_{\mathcal{B}:\mathcal{D}}$. We first prove that $\mathcal{N}_0 = \{f \in \mathcal{B}\langle\mathcal{X}\rangle : \ \mu(f^*f) = 0\}$ is a left ideal of $\mathcal{B}\langle\mathcal{X}\rangle$. It suffices to prove that if $f \in \mathcal{N}_0$ then $\mathcal{X} \cdot f \in \mathcal{N}_0$ and $b \cdot f \in \mathcal{N}_0$ for all $b \in \mathcal{B}$.

Since $b^*b \leq \|b^*b\|$ (in the C*-algebra $\mathcal{B}$), the positivity of μ implies

$$\mu(f^*(\|b^*b\| - b^*b)f) \geq 0,$$

that is $\mu(f^*b^*bf) \leq \|b^*b\|\mu(f^*f) = 0$, hence $b \cdot f \in \mathcal{N}_0$.

For $g = b_0\mathcal{X}b_1\mathcal{X} \cdots \mathcal{X}b_n$ a monomial in $\mathcal{B}\langle\mathcal{X}\rangle$, define $\mathfrak{p}(g) = M^n\|b_0\|\|b_1\| \cdots \|b_n\|$ (in particular, condition (2) states that $\|\mu(g)\| \leq \mathfrak{p}(g)$). Consider

$$\mathcal{B}\langle\langle\mathcal{X}\rangle\rangle_\mu = \{\sum_{n=0}^{\infty} f_n : f_n = \text{monomials in } \mathcal{B}\langle\mathcal{X}\rangle \text{such that } \sum_{n=0}^{\infty} \mathfrak{p}(f_n) < \infty\}.$$

$\mathcal{B}\langle\langle\mathcal{X}\rangle\rangle_\mu$ is a $*$-algebra (with the structure inherited from $\mathcal{B}\langle\mathcal{X}\rangle$) and μ extends to a positive map $\widetilde{\mu} : \mathcal{B}\langle\langle\mathcal{X}\rangle\rangle_\mu \longrightarrow \mathcal{D}$ via $\widetilde{\mu}(\sum_{n=0}^{\infty} f_n) = \sum_{n=0}^{\infty} \mu(f_n)$.

Take now $g_n = (2n)!\left[(1-2n)(n!)^2(4M)^{2n}\right]^{-1}\mathcal{X}^{2n}$, $n \geq 0$. Then $g_n = g_n^*$ and $\mathfrak{p}(g_n) \leq 4^{-n}$, so $g = \sum_{n=0}^{\infty} g_n$ is also a selfadjoint element of $\mathcal{B}\langle\langle\mathcal{X}\rangle\rangle_\mu$.

Since $g^2 = 1 - \left[(2M)^{-1}\mathcal{X}\right]^2$, we have that

$$\begin{aligned} 0 \leq \widetilde{\mu}(f^*g^*gf) &= \widetilde{\mu}\left(f^*[1 - (2M)^{-2}\mathcal{X}^2] \cdot f\right) \\ &= \mu(f^*f) - (2M)^{-2}\mu\left((\mathcal{X}f)^*\mathcal{X}f\right) \end{aligned}$$

hence $\mu\left((\mathcal{X}f)^*\mathcal{X}f\right) \leq 4M^2\mu(f^*f)$, so $\mathcal{X}f \in \mathcal{N}_0$.

Consider now $\mathcal{K} = \mathcal{B}\langle\mathcal{X}\rangle \otimes_{\mathcal{B}} \mathcal{D}$ with the right $\mathcal{D}$-module structure given by $(f(\mathcal{X}) \otimes d)d_1 = f(\mathcal{X}) \otimes dd_1$; also, since μ satisfies the condition (1), we can define a $\mathcal{D}$-valued sesquilinear inner-product structure on $\mathcal{K}$ (see [13], page 40) via

$$\langle f(\mathcal{X}) \otimes d_1, g(\mathcal{X}) \otimes d_2\rangle = d_2^*\mu(g(\mathcal{X})^*f(\mathcal{X}))d_1.$$

Let now $\mathcal{N} = \{\eta \in \mathcal{K} \colon \langle \eta, \eta \rangle = 0\}$. From the above argument on $\mathcal{N}_0$, we have that $\mathcal{N}$ is a left $\mathcal{B}\langle \mathcal{X} \rangle$-module. Finally, take E the completion of $\mathcal{K}/\mathcal{N}$ in the norm induced by the inner-product structure (see [13]) and let $\xi = 1 \otimes 1 + \mathcal{N}$. The multipliers with polynomials from $\mathcal{B}\langle \mathcal{X} \rangle$ are in the C*-algebra of bounded maps from $\mathcal{L}(E)$, since condition (2) ensures the boundness. Moreover, if $\phi(\cdot) = \langle \cdot \xi, \xi \rangle$ and a is the right multiplier with $\mathcal{X}$, then $\mu = \phi_a$.

Condition (3) is implied by the equality $\|\mathrm{Id}_m \otimes a\| = \|a\|$, where the first norm is in the C*-algebra $M_m(A)$ and the second norm in the C*-algebra A.

□

2. Infinite divisibility: the boolean case

The main result of this section is the non-commutative analogue of the following theorem (see [23]):

Theorem 2.1. *Any compactly supported real measure is infinitely divisible with respect to boolean convolution.*

We will use the following notion of boolean independence over a C*-algebra $\mathcal{B}$ (see [18]):

Definition 2.2. *Let B be a unital C*-algebra, $B \subseteq D$, $B \subseteq A$ be unital inclusions of $*$-algebras and $\phi : A \longrightarrow D$ be a unital B-bimodule map. A family $\{a_i\}_{i \in I}$ of selfadjoint elements from A is said to be* boolean independent *with respect to ϕ if*

$$\phi(A_1 A_2 A_3 \cdots) = \phi(A_1)\phi(A_2)\phi(A_3)\cdots$$

for all $A_k \in \mathcal{B}\langle a_{\epsilon(k)} \rangle_0$ (the $*$-algebra spanned by non-commutative polynomials in $a_{\epsilon(k)}$ and coefficients in $\mathcal{B}$ without a free term), with $\epsilon(k) \in I$ and $\epsilon(k) \neq \epsilon(k+1)$.

Let now $N \in \mathbb{N}$ and $\{\mu_j\}_{j=1}^N$ be a family of elements from $\Sigma_{\mathcal{B}:\mathcal{D}}$. We define their additive boolean convolution as follows. Consider the symbols $\{X_j\}_{j=1}^N$ such that $\mu_{X_j} : \mathcal{B}\langle X_j \rangle \cong \mathcal{B}\langle \mathcal{X} \rangle \longrightarrow \mathcal{B}$ coincides to μ_j. Then consider the $*$-algebra $\mathcal{B}\langle X_1, X_2, \dots, X_N \rangle$ with the conditional expectation μ such that its restrictions to $\mathcal{B}\langle X_j \rangle$ coincide to μ_j and the mixed moments of $X_1, X_2, \dots, X_N$ are calculated via the rule from Definition 2.2. The additive boolean convolution of $\{\mu_j\}_{j=1}^N$ is the unital $\mathcal{B}$-bimodule map

$$\uplus_{j=1}^N \mu_j = \mu_{X_1 + X_2 + \cdots + X_N} : \mathcal{B}\langle X_1 + X_2 + \cdots + X_N \rangle \cong \mathcal{B}\langle \mathcal{X} \rangle \longrightarrow \mathcal{D}$$

Remark 2.3. *If $\mu_j \in \Sigma_{\mathcal{B}:\mathcal{D}}$ for al $j = 1, \dots, N$, then $\uplus_{j=1}^N \mu_j$ is also in $\Sigma_{\mathcal{B}:\mathcal{D}}$.*

Proof. For each j, consider the right-$\mathcal{D}$ module $\mathcal{K}_j = \mathcal{B}\langle X_j \rangle \otimes_{\mathcal{B}} \mathcal{D}$ as in Proposition 1.2.

Note that the $\mathcal{D}$-submodule $1 \otimes \mathcal{D}$ is complemented in each $\mathcal{K}_j$, since, for all $f(X) \in \mathcal{B}\langle \mathcal{X} \rangle$, we have that $1 \otimes \mu_j(f(X_j)) \in 1 \otimes_{\mathcal{D}}$ and

$$\langle 1 \otimes 1, f(X_j) \otimes 1 - 1 \otimes \mu_j(f(X_j)) \rangle_j = \mu_j(f(X_j)^*) - \mu_j(f(X_j))^* = 0$$

Denote by $\mathcal{K}_j^0$ the complement of $1 \otimes_{\mathcal{D}}$ in $\mathcal{K}_j$ and let

$$\mathcal{K} = (1 \otimes \mathcal{D}) \oplus \bigoplus_{j=1}^N \mathcal{K}_j^0.$$

On $\mathcal{L}(\mathcal{K})$ we consider the map $\phi : T \mapsto \langle T 1 \otimes 1, 1 \otimes 1 \rangle$.

$\mathcal{B}\langle X_j\rangle$ can be seen as a algebra of linear maps on $\mathcal{K}_j$ via

$$f(X_j)\big[g(X_j)\otimes d\big] = \big(f(X_j)g(X_j)\big)\otimes d.$$

Since $f(X_j)^*$ is adjoint to $f(X_j)$ we have $\mathcal{B}\langle X_j\rangle \in \mathcal{L}(\mathcal{K})$. moreover, by setting the restrictions of $\mathcal{B}\langle X_j\rangle_0$ to each $\mathcal{K}_l^0$ to the 0 if $l \neq j$, we can see $\mathcal{B}\langle X_j\rangle$ as a subalgebra of $\mathcal{L}(\mathcal{K})$. Note that $\phi_{|\mathcal{B}\langle X_j\rangle_0} = \mu_j$.

If $j \neq l$ and $A_1 \in \mathcal{B}\langle X_j\rangle_0$ while $A_2 \in \mathcal{B}\langle X_l\rangle_0$, then

$$\begin{aligned} A_1A_2(1\otimes 1) &= A_1[\phi(A_2)(1\otimes 1)+\xi], \text{ where } \xi = A_2(1\otimes 1) - \phi(A_2)1\otimes 1 \in \mathcal{K}_l^0 \\ &= A_1\phi(A_2)(1\otimes 1). \end{aligned}$$

Iterating, we obtain that for $A_k \in \mathcal{B}\langle X_{\epsilon(k)}\rangle_0$ with $\epsilon(k) \neq \epsilon(k+1)$

$$\phi(A_1\cdots A_m) = \phi(A_1\phi(A_2)\cdots\phi(A_m)) = \phi(A_1)\cdots\phi(A_m).$$

that is X_j's are boolean independent in $\mathcal{L}(\mathcal{K})$ with respect to ϕ, and since the restrictions of ϕ to $\mathcal{B}\langle X_j\rangle$ are μ_j, we have that $\uplus_{j=1}^N \mu_j = \phi_{|\mathcal{B}\langle X_1+\cdots+X_N\rangle}$ so q.e.d.. ☐

Definition 2.4. *An element $\mu \in \Sigma_{\mathcal{B}:\mathcal{D}}$ is said to be $\biguplus$-infinite divisible if for any positive integer n there exist some $\mu_n \in \Sigma_{\mathcal{B}:\mathcal{D}}$ such that μ is the additive boolean convolutions of n copies of μ_n.*

The main result of this section si the following

Theorem 2.5. *Any element μ from $\Sigma_{\mathcal{B}:\mathcal{D}}$ is boolean infinitely divisible.*

To prove 2.5, we will need the following generalization of the notion of scalar boolean cumulants from [23]:

Definition 2.6. *In the above setting, let X be a selfadjoint element from $\mathcal{A}$. The* boolean cumulants *of X are defined as the multilinear maps $\{B_{n,X}\}_{n\geq 1}$, with $B_{n,X} : \mathcal{B}^n \longrightarrow \mathcal{D}$, given by the following recurrence:*

$$\phi(Xb_nXb_{n-1}\cdots Xb_1) = \sum_{k=1}^{n} \phi(Xb_n\cdots Xb_{k+1})B_{k,X}(b_k,\cdots,b_1).$$

As shown in [18], the $\mathcal{D}$-valued boolean cumulants defined above have the same additivity property as their scalar analogues (see [18], Corollary 4.6):

Proposition 2.7. *If, in the above setting, X and Y are boolean independent over ϕ, then, for all positive integers n, we have that*

$$B_{n,X+Y} = B_{n,X} + B_{n,Y}.$$

Definition 2.4 can be reformulated in terms of Proposition 2.7. Namely, for $\mu \in \Sigma_{\mathcal{B}:\mathcal{D}}$ we define the n-th boolean cumulant of μ as the multilinear map $B_{n,\mu} : \mathcal{B}^n \longrightarrow \mathcal{D}$ given by the recurrence:

$$\mu(\mathcal{X}b_1\mathcal{X}b_2\cdots\mathcal{X}b_n) = \sum_{k=1}^{n} B_{k,\mu}(b_1,\ldots,b_k)\mu(\mathcal{X}b_{k+1}\cdots\mathcal{X}b_n).$$

From Proposition 2.7, we have the following

Remark 2.8. *An element $\mu \in \Sigma_{\mathcal{B}:\mathcal{D}}$ is $\biguplus$-infinite divisible if for any positive integer n there exist $\mu_n \in \Sigma_{\mathcal{B}:\mathcal{D}}$ such that for all positive integers m we have that*

$$B_{m,\mu} = nB_{m,\mu_n}.$$

Before proving the main result, i. e. Theorem 2.5, we will first need the following result.

Lemma 2.9. *Let $B \subseteq A$ be a unital inclusion of C^*-algebras, $\mathcal{H}$ be a semi-inner product A-bimodule which is also a left B-module and Ω a symbol that commutes with A such that $\langle \Omega, \Omega \rangle = 1$ and $\langle \cdot\Omega, \Omega \rangle : \mathcal{L}(\mathcal{H} \oplus \Omega A) \longrightarrow A$ is a B-bimodule map.*

Let $T, \Lambda \in \mathcal{L}(\mathcal{H} \oplus \Omega A)$ be selfdajoint operators such that $T(\Omega A) = 0, T(\mathcal{H}) \subseteq \mathcal{H}$, $\Lambda(\mathcal{H}) = 0$, $\Lambda(\Omega A) \subseteq \Omega A$. For $\xi \in \mathcal{H}$ define the operators $a_\xi, a_\xi^* \in \mathcal{L}(\mathcal{H} + \oplus \Omega A)$ given by

$$\begin{cases} a_\xi \alpha = 0 & \alpha \in A \\ a_\xi \eta = \langle \eta, \xi \rangle & \eta \in \mathcal{H} \end{cases} \qquad\qquad \begin{cases} a_\xi^* \alpha = \xi\alpha & \alpha \in A \\ a_\xi^* \eta = 0 & \eta \in \mathcal{H} \end{cases}.$$

Then a_ξ, a_ξ^* are adjoint to each other and the boolean cumulants $\{B_{n,V}\}_n$ of $V = a_\xi + a_\xi^* + T + \Lambda$ with respect to the map $\langle \cdot\Omega, \Omega \rangle$ are given by:

$$\begin{array}{ll} (i) & B_{1,V}(b_1) = \langle \Lambda b_1 \Omega, \Omega \rangle \\ (ii) & B_{n,V}(b_n, \dots, b_1) = \langle a_\xi b_n T b_{n-1} \cdots T b_2 a_\xi^* b_1 \Omega, \Omega \rangle \text{ if } n \geq 2. \end{array}$$

Proof. The fact that a_ξ and a_ξ^* are adjoint to each other is just a trivial computation. For (i), note that $b_1\Omega \in \Omega A$, hence $\langle V b_1 \Omega, \Omega \rangle = \langle \Lambda b_1 \Omega, \Omega \rangle$.

For (ii), remark first that

$$a_\xi b_n T b_{n-1} \cdots T b_2 a_\xi^* b_1 \Omega \subset \Omega A$$

for all $b_1, \dots, b_n \in B$, since $a_\xi^* b_1 \Omega = \xi b_1 \in \mathcal{H}$, also $b_n T b_{n-1} \cdots T b_2 \mathcal{H} \subseteq \mathcal{H}$ and $a_\xi \mathcal{H} \subseteq \Omega A$.

We have that

$$\langle V b_n \cdots V b_1 \Omega, \Omega \rangle = \sum_{V_j \in \{a_\xi, a_\xi^*, T, \Lambda\}} \langle V_n b_n \cdots V_1 b_1 \Omega, \Omega \rangle.$$

Let us suppose that the term $\langle V_n b_n \cdots V_1 b_1 \Omega, \Omega \rangle$ does not cancel. Since $a_\xi(\Omega A) = T(\Omega A) = 0$, it follows that $V_1 \in \{a_\xi^*, \Lambda\}$. If $V_1 = a_\xi^*$, then $V_1 b_1 \Omega \in \mathcal{H}$ and, since

$$a_\xi^* \mathcal{H} = \Lambda \mathcal{H} = 0$$

it follows that $V_2 \in \{a_\xi, T\}$.

Also, since $T\mathcal{H} \subseteq \mathcal{H}$, if $V_2 = V_3 = \cdots = V_p = T$, then $V_{p+1} \in \{a_\xi, T\}$. Finally, note that $T b_n \cdots T b_2 a^* \xi b_1 \Omega \in \mathcal{H}$, henceforth

$$\begin{aligned} \langle V b_n \cdots V b_1 \Omega, \Omega \rangle &= \langle V b_n \cdots V b_2 \Lambda b_1 \Omega, \Omega \rangle \\ &\quad + \sum_{p=2}^{n} \langle V b_n \cdots V b_{p+1} a_\xi b_p T b_{p-1} \cdots T b_2 a^* \xi b_1 \Omega, \Omega \rangle \\ &= \langle V b_n \cdots V b_2 \Omega, \Omega \rangle \cdot \langle \Lambda b_1 \Omega, \Omega \rangle \\ &\quad + \sum_{p=2}^{n} \langle V b_n \cdots V b_{p+1} \Omega, \Omega \rangle \cdot \langle a_\xi b_p T b_{p-1} \cdots T b_2 a^* \xi b_1 \Omega, \Omega \rangle \end{aligned}$$

and the conclusion follows from (i) and Definition 2.6. □

Proof of the Theorem 2.5

Consider $\mathcal{K} = \mathcal{B}\langle\mathcal{X}\rangle \otimes_{\mathcal{B}} \mathcal{D}$ as for Remark 2.3. Denote by $\mathcal{K}^0$ the complement of $1 \otimes \mathcal{D}$ in $\mathcal{K}$ and let $\xi = \mathcal{X} \otimes 1 - 1 \otimes \mu(\mathcal{X}) \in \mathcal{K}^0$. Define the operator $a_\xi : \mathcal{K} \longrightarrow \mathcal{K}$ by

$$\begin{aligned} a_\xi(1 \otimes 1) &= 0 \\ a_\xi \eta &= 1 \otimes \langle \eta, \xi \rangle \text{ for } \eta \in \mathcal{K}^0 \end{aligned}$$

Remark that a_ξ is $\mathcal{D}$-linear, adjointable, and its adjoint is given by

$$\begin{aligned} a_\xi^*(1 \otimes 1) &= \xi 1 \\ a_\xi^* \eta &= 0 \text{ for } \eta \in \mathcal{K}^0 \end{aligned}$$

Let also let $\mathfrak{L}_{\mathcal{X}} : \mathcal{K} \longrightarrow \mathcal{K}$ be the selfadjoint map given by $f(\mathcal{X}) \otimes d \mapsto \mathcal{X} f(\mathcal{X}) \otimes d$. We will identify $\mathcal{B}$ with a subalgebra of $\mathcal{L}(\mathcal{K})$ via

$$b[f(\mathcal{X}) \otimes d] = [b f(\mathcal{X})] \otimes d.$$

Note that $\langle \cdot 1 \otimes 1, 1 \otimes 1 \rangle$ is a $\mathcal{B}$-bimodule map.

We have that

$$\begin{aligned} \langle \mathfrak{L}_{\mathcal{X}} b_1 \cdots \mathfrak{L}_{\mathcal{X}} b_n (1 \otimes 1), 1 \otimes 1 \rangle &= \langle \mathcal{X} b_1 \mathcal{X} b_2 \cdots \mathcal{X} b_n \otimes, 1 \otimes 1 \rangle \\ &= \mu(\mathcal{X} b_1 \mathcal{X} b_2 \cdots \mathcal{X} b_n) \end{aligned}$$

hence the distribution of $\mathfrak{L}_X$ with respect to $\langle \cdot 1 \otimes 1, 1 \otimes 1 \rangle$ coincides with μ.

Consider the selfadjoint map $\Lambda_\mu \in \mathcal{L}(\mathcal{K})$ given by

$$\begin{cases} \Lambda_\mu(1 \otimes d) = 1 \otimes \mu(\mathcal{X}) d, & d \in \mathcal{D} \\ \Lambda_\mu \eta = 0, & \eta \in \mathcal{K}^0 \end{cases}$$

and define $T \in \mathcal{L}(\mathcal{K})$ as the map $T = \mathfrak{L}_{\mathcal{X}} - (a_\xi + a_\xi^*) - \Lambda_\mu$. We have that T is selfadjoint, $T\mathcal{K}^0 \subseteq \mathcal{K}^0$ and $T(1 \otimes \mathcal{D}) = 0$.

Since $\mathfrak{L}_{\mathcal{X}} = a_\xi + a_\xi^* + \Lambda_\mu + T$ and its distribution with respect to $\langle \cdot 1 \otimes 1, 1 \otimes 1 \rangle$ is μ from Lemma 2.9 we have that the boolean cumulants of μ are given by

$$B_{1,\mu}(b_1) = \langle \Lambda_\mu b_1 (1 \otimes 1), 1 \otimes 1 \rangle \tag{4}$$

$$B_{n,\mu}(b_n, \dots, b_1) = \langle a_\xi b_n T b_{n-1} \cdots T b_2 a_\xi^* b_1 (1 \otimes 1), 1 \otimes 1 \rangle \text{ if } n \geq 2. \tag{5}$$

Fix N a positive integer. Let $\xi_N = \frac{1}{\sqrt{N}} \xi$, $\Lambda_N = \frac{1}{N} \Lambda_\mu$ and $Y_N \in \mathcal{L}(\mathcal{K})$ be the selfadjoint operator

$$Y_N = a_{\xi_N} + a_{\xi_N}^* + \Lambda_N + T.$$

Define $\mu_N \in \Sigma_{\mathcal{B}:\mathcal{D}}$ via

$$\mu_N(f(X)) = \langle f(Y_N)(1 \otimes 1), 1 \otimes 1 \rangle.$$

From Lemma 2.9, the boolean cumulants of μ_N are given by:

$$\begin{aligned} B_{1,\mu_N}(b_1) &= \langle \Lambda_N b_1 (1 \otimes 1), 1 \otimes 1 \rangle = \frac{1}{N} \langle \Lambda_\mu b_1 (1 \otimes 1), 1 \otimes 1 \rangle \\ &= \frac{1}{N} B_{1,\mu}(b_1), \end{aligned}$$

$$\begin{aligned} B_{n,\mu_N}(b_1, \dots, b_n) &= \langle a_{\xi_N} b_1 T b_2 \cdots T b_{n-1} a_{\xi_N}^* b_n (1 \otimes 1), 1 \otimes 1 \rangle \\ &= \frac{1}{N} \langle a_\xi b_1 T b_2 \cdots T b_{n-1} a_\xi^* b_n (1 \otimes 1), 1 \otimes 1 \rangle \\ &= \frac{1}{N} B_{n,\mu}(b_1, \dots, b_n) \end{aligned}$$

and the conclusion follows from Remark 2.8.

□

3. Infinite divisibility: the Free case

3.1. Preliminaries.

Definition 3.1. (see [27]) *Let $\mathcal{B}$ be a unital C*-algebra, $\mathcal{B} \subseteq \mathcal{A}$ be a unital inclusion of $*$-algebras and $\phi : \mathcal{A} \longrightarrow \mathcal{B}$ be a positive conditional expectation. A family $\{X_i\}_{i\in I}$ of selfadjoint elements from $\mathcal{A}$ is said to be* free *with respect to if ϕ if*

$$\phi(A_1 A_2 \cdots A_n) = 0$$

whenever $A_j \in \mathcal{B}\langle X_{\epsilon(j)}\rangle$ with $\epsilon(k) \neq \epsilon(k+1)$ and $\phi(A_j) = 0$.

In the above setting, let X be a selfadjoint element from $\mathcal{A}$. The free cumulants of X are the multilinear functions $\kappa_{n,X} : \mathcal{B}^n \longrightarrow \mathcal{B}$ given by the recurrence:

$$\begin{aligned}\phi(Xb_1Xb_2\cdots Xb_n) \quad = \\ \sum_{p=1}^{n} \sum_{1<j_1<\cdots<j_{p-1}} \kappa_{p,X}((b_1\phi(Xb_2\cdots Xb_{j-1}), (b_{j_1+1}\phi(Xb_{j_1+2}\cdots Xb_{j_2}), \dots \\ \dots, (b_{j_{p_1}}\phi(Xb_{j_{p-1}+1}\cdots Xb_n)).\end{aligned}$$

More intuitively, the above relation can be graphically illustrated by the picture below, where the boxes stand for the application of ϕ and the oblique lines signify that each $Y_s = \phi(Xb_{j_s+2}\cdots Xb_{j_{s+1}})$ are multiplied with b_{j_s+1} in the arguments of the free cumulants:



The free cumulants have the following additivity property (see [24], [17]):

Proposition 3.2. *If X, Y are free in the sense of Definition 3.1, then*

$$\kappa_{n,X+Y} = \kappa_{n,X} + \kappa_{n,Y}.$$

Let now $N \in \mathbb{N}$ and $\{\mu_j\}_{j=1}^N$ be a family of elements from $\Sigma_{\mathcal{B}}$. We define their additive free convolution similarly to be boolean case: Consider the symbols $\{X_j\}_{j=1}^N$; on the algebra $\mathcal{B}\langle X_1, X_2, \dots, X_N\rangle$ take the conditional expectation μ such that $\mu \circ \tau_{X_j} = \mu_j$ and the mixed moments of $X_1, \dots, X_n$ are computed via the rules from Definition 3.1. The free additive convolution of $\{\mu_j\}_{j=1}^N$ is the conditional expectation

$$\boxplus_{j=1}^N \mu_j = \mu \circ \tau_{X_1+X_2+\cdots+X_N} : \mathcal{B}\langle X_1 + X_2 + \cdots + X_N\rangle \cong \mathcal{B}\langle \mathcal{X}\rangle \longrightarrow \mathcal{B}.$$

We have that $\boxplus_{j=1}^N \mu_j$ is also an element of $\Sigma_{\mathcal{B}}$: in [24], Theorem 3.5.6, it is shown that μ, defined as above, is a positive conditional expectation, therefore so is $\mu \circ \tau_{X_1+X_2+\cdots+X_N}$.

Definition 3.3. *An element $\mu \in \Sigma_{\mathcal{B}}$ is said to be $\boxplus$-infinite divisible if for any positive integer N there exists $\mu_N \in \Sigma_{\mathcal{B}}$ such that μ is the free additive convolution of N copies of μ_N.*

3.2. Free cumulants and $\boxplus$-infinite divisibility.

Definition 3.4. *Let $\mu \in \Sigma_{\mathcal{B}}$. Using the relations from Section 3.1, we define the free cumulants of μ as the multilinear maps $\kappa_{n,\mu} : \mathcal{B}^n \longrightarrow \mathcal{B}$ given by the recurrence:*

$$\mu(\mathcal{X}b_1\mathcal{X}b_2\cdots\mathcal{X}b_n) = \sum_{p=1}^{n} \sum_{1<j_1<\cdots<j_{p-1}} \kappa_{p,\mu}((b_1\mu(\mathcal{X}b_2\cdots\mathcal{X}b_{j-1}), (b_{j_1+1}\mu(\mathcal{X}b_{j_1+2}\cdots\mathcal{X}b_{j_2}), \dots$$
$$\dots, (b_{j_{p1}}\mu(\mathcal{X}b_{j_{p-1}+1}\cdots\mathcal{X}b_n)).$$

Remark 3.5. *Using Proposition 3.2, we can reformulate Definition 3.4 in terms of free cumulants. More precisely, $\mu \in \Sigma_{\mathcal{B}}$ is $\boxplus$-infinitely divisible if for any positive integer N there exists some $\mu_N \in \Sigma_{\mathcal{B}}$ such that for all m*

$$\kappa_{m,\mu} = N \cdot \kappa_{m,\mu_N}.$$

For $\mu \in \Sigma_{\mathcal{B}}$, define the conditional expectation $\rho_\mu : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{B}$ generated by

$$\rho_\mu(\mathcal{X}b_1\mathcal{X}b_2\cdots\mathcal{X}b_n) = \kappa_{n,\mu}(b_1, b_2, \cdots, b_n).$$

Proposition 3.6. *If $\mu \in \Sigma_{\mathcal{B}}$ is $\boxplus$-infinitely divisible then the restriction of ρ_μ to $\mathcal{B}\langle\mathcal{X}\rangle_0$ is positive.*

Proof. Fix N and suppose that μ is the free additive convolution of N copies of μ_N. Note that, for $n \leq 1$,

$$\mu_N(\mathcal{X}b_1\cdots\mathcal{X}b_n) = \frac{1}{N}\kappa_{n,\mu}(b_1, \dots, b_n) + O(\frac{1}{N^2}). \tag{6}$$

The assertion is trivial for $n = 1$. Suppose that (6) is true for $n < m$. Since the free cumulants are multilinear, for all $1 = l_1 < l_2 < \cdots < l_{p+1} = m + 1$ and $Y_s = b_{l_s}\mu_N(\mathcal{X}b_{l_s+1}\cdots\mathcal{X}b_{l_{s+1}-1})$, $(1 \leq s \leq p)$ we have that

$$Y_s = \begin{cases} b_{l_s} & \text{if } l_{s+1} = l_s + 1 \\ O(\frac{1}{N}) & \text{if } l_{s+1} = l_s + 1 \end{cases}$$

hence

$\kappa_{p,\mu_N}(Y_1, \dots, Y_p) = \frac{1}{N}\kappa_{p,\mu}(Y_1, \dots, Y_p) = O(\frac{1}{N^2})$ unless $l_{s+1} = l_s + 1$ for all $s \in \{1, \dots, p\}$, i. e. $p = m$.

Definition 3.4 gives

$$\mu_N(\mathcal{X}b_1\cdots\mathcal{X}b_m) = \kappa_{m,\mu_N}(b_1, \dots, b_m) + O(\frac{1}{N^2}) = \frac{1}{N}\kappa_{m,\mu}(b_1, \dots, b_m)O(\frac{1}{N^2})$$

that is (6). It follows that

$$\lim_{N\longrightarrow\infty} N \cdot \mu_N(\mathcal{X}b_1\cdots\mathcal{X}b_n) = \kappa_{n,\mu}(b_1, \dots, b_n) \tag{7}$$

Fix now $f(\mathcal{X}) \in \mathcal{B}\langle\mathcal{X}\rangle_0$. Then

$$\rho_\mu(f(\mathcal{X})^*f(\mathcal{X})) = \lim_{N\longrightarrow\infty}[N\mu_N(f(\mathcal{X})^*f(\mathcal{X}))] \geq 0 \text{ since all } \mu_N \text{ are positive.}$$

□

Lemma 3.7. *Let A be a unital C^*-algebra and $\mathcal{H}$ a semi-inner-product A-module which is also a left A-module. Let $\mathcal{T}(\mathcal{H})$ be the* full Fock A-module *over $\mathcal{H}$, that is*

$$\mathcal{T}(\mathcal{H}) = A \oplus \mathcal{H} \oplus (\mathcal{H} \otimes_B \mathcal{H}) \oplus (\mathcal{H} \otimes_B \mathcal{H} \otimes_B \mathcal{H}) \oplus \dots.$$

Fix $\xi \in \mathcal{H}$, $\beta \in A$, $T \in \mathcal{L}(\mathcal{H})$, T and β selfadjoint, and define the maps ($\alpha \in A$, $\eta_1, \dots, \eta_n \in \mathcal{H}$)

$$\begin{cases} a_\xi \alpha = 0 \\ a_\xi \eta_1 \otimes \eta_2 \otimes \cdots \otimes \eta_n = \langle \eta, \xi \rangle \eta_2 \otimes \cdots \otimes \eta_n \end{cases}$$

$$\begin{cases} a_\xi^* \alpha = \xi \alpha \\ a_\xi^* \eta_1 \otimes \cdots \otimes \eta_n = \xi \otimes \eta_1 \otimes \cdots \otimes \eta_n \end{cases}$$

$$\begin{cases} p(T)\alpha = 0 \\ p(T)\eta_1 \otimes \eta_2 \otimes \cdots \otimes \eta_n = T(\eta_1) \otimes \eta_2 \otimes \cdots \otimes \eta_n \end{cases}$$

Then a_ξ, a_ξ^* are adjoint to each other, $p(T)$ is selfadjoint and the cumulants of $X = a_\xi + a_\xi^* + \widetilde{T} + \beta\chi_A$ with respect to the conditional expectation $\langle \cdot 1, 1\rangle$ are given by

$$\begin{aligned} \kappa_{1,X}(b_1) &= \beta b_1 \\ \kappa_{n,X}(b_1, \dots, b_n) &= \langle a_\xi b_1 p(T) b_2 \cdots p(T) b_{n-1} a^* \xi b_1 1, 1\rangle. \end{aligned}$$

Proof. For $\kappa_{1,X}$ the assertion is trivial. To prove the relation for higher order free cumulants, let us fix $N > 1$ and consider $\{(\mathcal{H}_{j,N}, T_{j,N}, \xi_{j,N})\}_{j=1}^N$ to be a set of N identical copies of $(\mathcal{H}, T, (\frac{1}{\sqrt{N}})\xi)$ from above. Let

$$\begin{aligned} \mathcal{H}_N &= \mathcal{H}_{1,N} \oplus \mathcal{H}_{2,N} \oplus \cdots \oplus \mathcal{H}_{N,N} \\ \xi_N &= \xi_{1,N} \oplus \xi_{2,N} \oplus \cdots \oplus \xi_{N,N} \in \mathcal{H}_N \\ T_N &= T_{1,N} \oplus \cdots \oplus T_{N,N} \in \mathcal{L}(\mathcal{H}_N) \\ X_N &= a_{\xi_N} + a_{\xi_N}^* + P(T_N) + \beta\chi_A \in \mathcal{L}(\mathcal{H}_N) \\ X_{j,N} &= a_{\xi_{j,N}} + a_{\xi_{j,N}^*} + p(T_{j,N}) + \frac{1}{N}\beta\chi_A \in \mathcal{L}(\mathcal{H}_N). \end{aligned}$$

First, note that X and X_N are identically distributed with respect to $\langle \cdot 1, 1\rangle$. Then $X_N = X_{1,N} + \cdots + X_{N,N}$, $X_{j,N}$ are identically distributed and free. To see that, consider $A_1, \dots, A_m$, $A_k \in A\langle X_{\epsilon(k),N}\rangle$ with $\langle A_k 1, 1\rangle = 0$ and $\epsilon(k) \neq \epsilon(k+1)$. We have to show that $\langle A_m \cdots A_1 1, 1\rangle = 0$.

Since $\langle A_1 1, 1\rangle = 0$, we have that $A_1 1 \in \mathcal{T}(\mathcal{H}_{\epsilon(1),N}) \ominus A$. Also, $\langle A_2 1, 1\rangle = 0$, so $A_2 A_1 1 \in (\mathcal{T}(\mathcal{H}_{\epsilon(2),N}) \ominus A) \otimes (\mathcal{T}(\mathcal{H}_{\epsilon(1),N}) \ominus A)$. Iterating, we obtain

$$A_m \cdots A_2 A_1 1 \in (\mathcal{T}(\mathcal{H}_{\epsilon(m),N} \ominus A) \otimes \cdots \otimes (\mathcal{T}(\mathcal{H}_{\epsilon(1),N}) \ominus A)$$

so $\langle A_m \cdots A_2 A_1 1, 1\rangle = 0$.

Using (7) and denoting $\mathcal{V} = \{\frac{1}{\sqrt{N}} a_\xi, \frac{1}{\sqrt{N}} a_\xi^*, p(T), \frac{1}{N}\beta\chi_A\}$ we have ($n \geq 2$):

$$\begin{aligned} \kappa_{n,X}(b_1, \dots, b_n) &= \kappa_{n,X_N}(b_1, \dots, b_n) \\ &= \lim_{N \longrightarrow \infty} N\langle X_N b_1 \cdots X_N b_n 1, 1\rangle \\ &= \sum_{V_i \in \mathcal{V}} \langle V_1 b_1 \cdots V_n b_n 1, 1\rangle \end{aligned}$$

But $\langle V_1 b_1 \cdots V_n b_n 1, 1\rangle = 0$ unless $V_n \in \{\frac{1}{\sqrt{N}} a_\xi^*, \frac{1}{N}\beta\chi_A\}$ and $V_1 \in \{\frac{1}{\sqrt{N}} a_\xi, \frac{1}{N}\beta\chi_A\}$, therefore

$$\langle X_N b_1 \cdots X_N b_n 1, 1\rangle = \langle \frac{1}{\sqrt{N}} a_\xi b_1 p(T) b_2 \cdots p(T) b_{n-1} \frac{1}{\sqrt{N}} a_\xi^* b_1 1, 1\rangle + O(\frac{1}{N^2})$$

hence the conclusion. □

Theorem 3.8. *The conditional expectation $\mu \in \Sigma_{\mathcal{B}}$ is $\boxplus$-infinitely divisible if and only if the restriction of ρ_μ to $\mathcal{B}\langle\mathcal{X}\rangle_0$ is positive.*

Proof. Suppose that $\rho_{\mu|\mathcal{B}\langle X\rangle_0}$ is positive. Then (see [13], pag. 42) $\mathcal{B}\langle\mathcal{X}\rangle_0$ is a semi-inner-product $\mathcal{B}$-module with respect to the pairing

$$\langle f(\mathcal{X}), g(\mathcal{X})\rangle = \rho_\mu(g(\mathcal{X})^* f(\mathcal{X})).$$

Consider the selfadjoint map $T : \mathcal{B}\langle\mathcal{X}\rangle_0 \longrightarrow \mathcal{B}\langle\mathcal{X}\rangle_0$ given by $T(f(\mathcal{X})) = \mathcal{X}f(\mathcal{X})$ and denote by V the map from $\mathcal{L}(\mathcal{T}(\mathcal{B}\langle\mathcal{X}\rangle_0))$ defined as

$$V = a_{\mathcal{X}} + a^*_{\mathcal{X}} + \widetilde{T} + \mu(\mathcal{X})\mathrm{Id}.$$

From Lemma 3.7 we have that the free cumulants of V with respect to $\langle \cdot 1, 1\rangle$ are given by

$$\begin{aligned}
\kappa_{1,V}(b_1) &= \mu(\mathcal{X})(b_1) = \kappa_{1,\mu}(b_1)\\
\kappa_{n,V}(b_1,\dots,b_n) &= \langle a_{\mathcal{X}} b_1 \widetilde{T} b_2 \cdots \widetilde{T} b_{n-1} a^*_{\mathcal{X}} b_n 1, 1\rangle = \langle b_1 \mathcal{X} b_2 \cdots \mathcal{X} b_n, \mathcal{X}\rangle\\
&= \kappa_\mu(b_1,\cdots,b_n)
\end{aligned}$$

Fix a positive integer N. From Remark 3.5 it suffices to find a selfadjoint element from $\mathcal{L}(\mathcal{T}(\mathcal{B}\langle\mathcal{X}\rangle_0))$ whose free cumulants are proportional to the free cumulants of V by a factor of $\frac{1}{N}$. Define

$$V_N = \frac{1}{\sqrt{N}} a_{\mathcal{X}} + \frac{1}{\sqrt{N}} a^*_{\mathcal{X}} + \widetilde{T} + \frac{1}{N}\mu(\mathcal{X})\mathrm{Id}.$$

Applying again Lemma 3.7, the free cumulants of V_N are

$$\begin{aligned}
\kappa_{1,V_N}(b_1) &= \frac{1}{N}\mu(\mathcal{X})(b_1) = \frac{1}{N}\kappa_{1,V}(b_1)\\
\kappa_{n,V_N}(b_1,\dots,b_n) &= \langle \frac{1}{\sqrt{N}} a_{\mathcal{X}} b_1 \widetilde{T} b_2 \cdots \widetilde{T} b_{n-1} \frac{1}{\sqrt{N}} a^*_{\mathcal{X}} b_n 1, 1\rangle\\
&= \frac{1}{N}\kappa_{n,V}(b_1,\cdots,b_n)
\end{aligned}$$

The converse implication is Proposition 3.6.

□

4. INFINITE DIVISIBILITY: THE C-FREE CASE

In this section we aim to extend the results from Section 3 to the case $\mu \in \Sigma_{\mathcal{B}:\mathcal{D}}$. First, we will need a suitable definition for the free additive convolution of elements from $\Sigma_{\mathcal{B}:\mathcal{D}}$; in this setting, if $\mathcal{B}$ is simply replaced by $\mathcal{D}$ in Definition 3.1, the resulting relation does not uniquely determine the joint moments of $X_1,\dots,X_n$. As shown in [5], [8], a more suitable approach is the conditional freeness (see also [17], [6]).

Definition 4.1. *Let $\mathcal{B}$ be a unital C*-algebra $\mathcal{B} \subseteq \mathcal{A}$, $\mathcal{B} \subseteq \mathcal{D}$ be unital inclusions of $*$-algebras, $\varphi : \mathcal{A} \longrightarrow \mathcal{B}$ be a conditional expectation and $\theta : \mathcal{A} \longrightarrow \mathcal{D}$ be a unital $\mathcal{B}$-bimodule map.*

The family $\{X_i\}_{i\in I}$ of selfadjoint elements from $\mathcal{A}$ is said to be c-free with respect to (θ, φ) if

- (i) *the family $\{X_i\}_{i\in I}$ is free with respect to φ*
- (ii) *$\theta(A_1 A_2 \cdots A_n) = \theta(A_1)\theta(A_2)\cdots\theta(A_n)$ for all $A_i \in \mathcal{B}\langle X_{\epsilon(i)}\rangle$ such that $\varphi(A_i) = 0$ and $\epsilon(k) \neq \epsilon(k+1)$.*

Let X be a selfadjoint element from $\mathcal{A}$. The c-free cumulants of X are the multilinear functions ${}^{c}\kappa_{n,X} : \mathcal{B}^n \longrightarrow \mathcal{D}$ given by the recurrence:

$$\begin{aligned}\theta(\mathcal{X}b_1\mathcal{X}b_2\cdots\mathcal{X}b_n) &= \sum_{p=1}^{n}\sum_{\substack{1<j_1<\cdots<j_p\\ l_p=n-1}} \theta(\mathcal{X}b_1\cdots\mathcal{X}b_{l_1})\cdot\\ &\qquad {}^{c}\kappa_{p,\mathcal{X}}\left(b_{l_1+1}\varphi(\mathcal{X}b_{l_1+2}\cdots\mathcal{X}b_{l_2}),\dots,b_{l_{p-1}+1}\varphi(\mathcal{X}b_{l_{p-1}+2}\cdots\mathcal{X}b_{l_p}),b_n\right)\end{aligned}$$

As in the previous section, the above equation can be represented more intuitively by the picture below, where the dark boxes stand for the application of θ, the light ones for the application of φ and the oblique lines signify that each $Y_s = \phi(\mathcal{X}b_{l_s+2}\cdots\mathcal{X}b_{l_{s+1}})$ are multiplied with b_{l_s+1} in the arguments of the c-free cumulants, except for b_n.



The c-free cumulants have the following additivity property (see [17], [15]):

Proposition 4.2. *If X, Y are c-free with respect to (θ, φ) in the sense of Definition 4.1, then*

$$\begin{aligned}\kappa_{n,X+Y} &= \kappa_{n,X} + \kappa_{n,Y}\\ {}^{c}\kappa_{n,X+Y} &= {}^{c}\kappa_{n,Y} + {}^{c}\kappa_{n,Y}\end{aligned}$$

where $\kappa_{n,X}$ is the n-th free cumulant of X with respect to the conditional expectation φ.

Let now N be a positive integer and $\{(\mu_i,\nu_i)\}_{i=1}^N$ be a family from $\Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$. We define their additive c-free convolution similarly to be boolean and free case: Consider the selfadjoint symbols $\{X_i\}_{i=1}^N$ and the mappings

$$\begin{aligned}\mu : \mathcal{B}\langle X_1, X_2, \dots, X_N\rangle &\longrightarrow \mathcal{D}\\ \nu : \mathcal{B}\langle X_1, X_2, \dots, X_N\rangle &\longrightarrow \mathcal{B}\end{aligned}$$

such that such that $\mu\circ\tau_{X_i} = \mu_i$ and $\nu\circ\tau_{X_i} = \nu_i$ for all $i = 1, \dots, N$ and the mixed moments of μ and ν are computed according to Definition 4.1. The c-free additive convolution of $\{(\mu_i,\nu_i)\}_{i=1}^N$ is the pair $(\mu_c,\nu_c) = \boxed{\mathrm{c}}_{i=1}^N(\mu_i,\nu_i)$, where

$$\begin{aligned}\nu_c &= \nu\circ\tau_{X_1+X_2+\cdots+X_N} = \boxplus_{i=1}^n \nu_i \in \Sigma_{\mathcal{B}}\\ \mu_c &= \mu\circ\tau_{X_1+X_2+\cdots+X_N} : \mathcal{B}\langle X_1+X_2+\cdots+X_N\rangle \cong \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}\end{aligned}$$

Definition 4.3. *A pair $(\mu,\nu) \in \Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$ is said to be $\boxed{c}$-infinite divisible if for any positive integer N there exists $(\mu_N,\nu_N) \in \Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$ such that (μ,ν) is the c-free additive convolution of N copies of (μ_N,ν_N).*

4.1. **C-free cumulants and infinite divisibility.**

Definition 4.4. *The c-free cumulants of the pair* $(\mu, \nu) \in \Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$ *are the multilinear functions* ${}^{c}\kappa_{n,\mu,\nu} : \mathcal{B}^n \longrightarrow \mathcal{D}$ *given by the recurrence:*

$$\begin{aligned}\mu(\mathcal{X}b_1\mathcal{X}b_2\cdots\mathcal{X}b_n) &= \sum_{p=1}^{n}\sum_{\substack{1<j_1<\cdots<j_p\\ l_p=n-1}} \mu(\mathcal{X}b_1\cdots\mathcal{X}b_{l_1})\cdot\\ &\quad {}^{c}\kappa_{p,\mu,\nu}\left(b_{l_1+1}\nu(\mathcal{X}b_{l_1+2}\cdots\mathcal{X}b_{l_2}),\dots,b_{l_{p-1}+1}\nu(\mathcal{X}b_{l_{p-1}+2}\cdots\mathcal{X}b_{l_p}),b_n\right)\end{aligned}$$

Remark 4.5. *As in the previous section, can reformulate Definition 4.4 in terms of free and c-free cumulants. More precisely, the pair* (μ, ν) *is* $\boxed{c}$*-infinitely divisible if for any positive integer* N *there exists some* $(\mu_N, \nu_N) \in \Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$ *such that for all* m *we have that* $\kappa_{m,\nu} = N\kappa_{m,\nu_N}$ *and* ${}^{c}\kappa_{m,\mu,\nu} = N^{c}\kappa_{m,\mu,\nu}$.

□

Define the map ${}^{c}\rho_{\mu,\nu} : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}$ as the $\mathcal{B}$-bimodule extension of

$${}^{c}\rho_{\mu,\nu}(\mathcal{X}b_1\cdots\mathcal{X}b_n) = {}^{c}\kappa_{n,\mu,\nu}(b_1,\dots,b_n).$$

Proposition 4.6. *Suppose that* $(\mu, \nu) \in \Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$ *is* $\boxed{c}$*-infinitely divisible. Then the restriction of* ${}^{c}\rho_{\mu,\nu}$ *to* $\mathcal{B}\langle\mathcal{X}\rangle_0$ *satisfies property (1)(see Introduction).*

Proof. Fix $N > 1$ and suppose that (μ, ν) is the c-free additive convolution of n copies of (μ_N, ν_N). As in the proof of Proposition 3.6, we will first show that

$$\mu_N(\mathcal{X}b_1\cdots\mathcal{X}b_n) = \frac{1}{N}{}^{c}\kappa_{n,\mu,\nu}(b_1,\dots,b_n) + O(\frac{1}{N^2}). \tag{8}$$

For $n = 1$ the assertion is trivial. Suppose that (8) holds true for $n < m$. Since the c-free cumulants of are multilinear, for all $1 = l_1 < l_2 < \cdots < l_{p+1} < m$ and $Y_s = b_{l_s}\nu_N(\mathcal{X}b_{l_s+1}\cdots\mathcal{X}b_{l_{s+1}-1})$, $(1 \le s \le p)$ we have that

$$Y_s = \begin{cases} b_{l_s} & \text{if } l_{s+1} = l_s + 1\\ O(\frac{1}{N}) & \text{if } l_{s+1} = l_s + 1\end{cases}$$

hence ${}^{c}\kappa_{p,\mu_N,\nu_N}(Y_1,\dots,Y_p) = \dfrac{1}{N}{}^{c}\kappa_{p,\mu,\nu}(Y_1,\dots,Y_p) = O(\dfrac{1}{N^2})$, unless $l_{s+1} = l_s + 1$ for all $s \in \{1,\dots,p\}$.

Definition 4.4 gives

$$\begin{aligned}\mu_N(\mathcal{X}b_1\cdots\mathcal{X}b_m) &= \kappa_{m,\mu_N,\nu_N}(b_1,\dots,b_m) +\\ &\quad \sum_{s=1}^{m-1}{}^{c}\kappa_{s,\mu_N,\nu_N}(b_1,\dots,b_s)\mu_N(\mathcal{X}b_{s+1}\ldots\mathcal{X}b_m) + O(\frac{1}{N^2})\end{aligned}$$

hence (8) follows from the induction hypothesis. Therefore

$$\lim_{N\longrightarrow\infty} N\mu_N(\mathcal{X}b_1\cdots\mathcal{X}b_n) = {}^{c}\kappa_{n,\mu,\nu}(b_1,\dots,b_n) \tag{9}$$

Fix now a family $\{f_i(\mathcal{X})\}_{i=1}^n$ in $\mathcal{B}\langle\mathcal{X}\rangle_0$. Then

$$[{}^{c}\rho_{\mu,\nu}(f_j(\mathcal{X})^*f_i(X))]_{i,j=1}^n = \lim_{N\longrightarrow\infty}[N\mu_N(f_j(\mathcal{X})^*f_i(X))]_{i,j=1}^n = \geq 0.$$

since each μ_N satisfies (1). □

Lemma 4.7. *Let $\mathcal{B} \subseteq \mathcal{D}$ be unital inclusion of C^*-algebras, $\mathcal{K}$ be a semi- inner-product $\mathcal{D}$-bimodule and $\mathcal{H}$ be a semi- inner-product $\mathcal{B}$-bimodule. Consider*

$$\mathcal{E} = (\mathcal{T}(\mathcal{H}) \otimes_{\mathcal{B}} \mathcal{D}) \oplus (1_{\mathcal{B}} \otimes_{\mathcal{B}} \Omega\mathcal{D}) \oplus (\mathcal{T}(\mathcal{H}) \otimes_{\mathcal{B}} \mathcal{K})\,.$$

Fix $\xi \in \mathcal{H}$, $\eta \in \mathcal{K}$ and $t \in \mathcal{L}(\mathcal{H})$, $T \in \mathcal{L}(\mathcal{K})$ selfadjoints. Define the maps a_ξ, a_ξ^*, A_η, A_η^*, $p(t)$, $P(T)$ from $\mathcal{L}(\mathcal{E})$ given by:

$$\begin{cases} a_{\xi|1_{\mathcal{B}}\otimes\Omega\mathcal{D}} \equiv 0 \\ a_\xi(f_1 \otimes \dots f_n) \otimes d = (\langle f_1, \xi\rangle f_2 \otimes \cdots \otimes f_n) \otimes d \\ a_\xi(f_1 \otimes \dots f_n) \otimes \eta = (\langle f_1, \xi\rangle f_2 \otimes \cdots \otimes f_n) \otimes \eta \end{cases}$$

$$\begin{cases} a^*_{\xi|1_{\mathcal{B}}\otimes\Omega\mathcal{D}} \equiv 0 \\ a^*_\xi(f_1 \otimes \dots f_n) \otimes d = (\langle f_1, \xi\rangle f_2 \otimes \cdots \otimes f_n) \otimes d \\ a^*_\xi(f_1 \otimes \dots f_n) \otimes \eta = (\langle f_1, \xi\rangle f_2 \otimes \cdots \otimes f_n) \otimes \eta \end{cases}$$

$$\begin{cases} p(t)_{|1_{\mathcal{B}}\otimes\Omega\mathcal{D}} \equiv 0 \\ p(t)(f_1 \otimes \dots f_n) \otimes d = (tf_1) \otimes f_2 \otimes \cdots \otimes f_n) \otimes d \\ p(t)(f_1 \otimes \dots f_n) \otimes \eta = (tf_1) \otimes f_2 \otimes \cdots \otimes f_n) \otimes \eta \end{cases}$$

$$\begin{cases} A_\eta(1 \otimes \zeta) = 1 \otimes \Omega\langle \zeta, \eta\rangle \\ A_{\eta|\mathcal{E}\ominus(1\otimes\mathcal{K})} \equiv 0 \end{cases} \qquad \begin{cases} A^*_\eta(1 \otimes \Omega d) = 1 \otimes \eta d \\ A^*_{\eta|\mathcal{E}\ominus(1\otimes\Omega\mathcal{D})} \equiv 0 \end{cases}$$

$$\begin{cases} P(T)(1 \otimes \zeta) = 1 \otimes T\zeta \\ P(T)_{|\mathcal{E}\ominus(1\otimes\mathcal{K})} \equiv 0 \end{cases}$$

Define also $I_1 = Id_{(\mathcal{T}(\mathcal{H})\otimes\mathcal{D})\oplus(\mathcal{T}(\mathcal{H})\otimes\mathcal{K})} \oplus 0 \in \mathcal{L}(\mathcal{E})$ and $I_2 = Id_{1\otimes\Omega\mathcal{D}} \oplus 0 \in \mathcal{L}(\mathcal{E})$. Consider $X = a_\xi + a_\xi^* + p(t) + \lambda_1 I_1 + A_\eta + A_\eta^* + \lambda_2 I_2$ where $\lambda_1 \in \mathcal{B}$, $\lambda_2 \in \mathcal{D}$ selfadjoint. Then the free and c-free cumulants of X with respect to $(\theta, \varphi) = (\langle \cdot 1 \otimes \Omega, 1 \otimes \Omega\rangle, \langle \cdot 1 \otimes 1, 1 \otimes 1\rangle)$ are given by the following relations:

$$\begin{cases} \kappa_{1,X}(b_1) = \lambda_1 b_1 \\ \kappa_{n,X}(b_1, \dots, b_n) = \langle a_\xi b_1 p(t) b_2 \cdots p(t) b_{n-1} a^*_\xi b_1 (1 \otimes 1), 1 \otimes 1\rangle \end{cases}$$

$$\begin{cases} {}^c\kappa_{1,X}(b_1) = \lambda_2 b_1 \\ {}^c\kappa_{n,X}(b_1, \dots, b_n) = \langle A_\eta b_1 P(T) b_2 \cdots P(T) b_{n-1} A^*_\eta b_1 (1 \otimes \Omega), 1 \otimes \Omega\rangle \end{cases}$$

Proof. The results are trivial for $\kappa_{1,X}$ and ${}^c\kappa_{1,X}$. For $\kappa_{n,X}$ note that $a_\xi, a_\xi^*, p(t)$ map $\mathcal{T}(\mathcal{H}) \otimes 1$ in $\mathcal{T}(\mathcal{H}) \otimes 1$ and, since $1 \otimes 1 \in \mathcal{T}(\mathcal{H}) \otimes 1$, the result reduces to Lemma **??**.

To prove the formula for ${}^c\kappa_{n,X}$, let us first note $V_0 = a_\xi + a_\xi^* + p(t) + \lambda_1 I_1$, $V_1 = A_\eta^*$, $V_2 = A_\eta$, $V_3 = P(T)$, $V_4 = \lambda_2 I_2$ and

$$J(n) = \{\vec{u} = (u_n, \dots, u_1) : 0 \leq u_k \leq 4\}.$$

Finally, for $b_1, \dots, b_n \in \mathcal{B}$, denote $\epsilon(\vec{u}) = \theta(V_{u_n} b_n \cdots V_{u_1} b_1)$.

Note that $V_{k|\mathcal{T}(\mathcal{H})\otimes 1} \equiv 0$ $(1 \leq k \leq 4)$ and $V_{0|1\otimes\Omega\mathcal{D}} \equiv 0$. The latest implies that $\epsilon(\vec{u}) = 0$ unless $V_{u_1} \in \{A_\eta^*, \lambda_2 I_2\}$, hence

$$\begin{aligned} \theta(X b_n \cdots X b_1) &= \sum_{\vec{u}\in J(n)} \epsilon(\vec{u}) = \sum_{\substack{\vec{u}\in J(n)\\ u_1=1}} \epsilon(\vec{u}) + \sum_{\substack{\vec{u}\in J(n)\\ u_1=4}} \epsilon(\vec{u}) \\ &= \sum_{\substack{\vec{u}\in J(n)\\ u_1=4}} \epsilon(\vec{u}) + \sum_{\vec{u}\in J(n-1)} \theta(V_{u_{n-1}} b_n \cdots V_{u_1} b_2)\lambda_2 \end{aligned}$$

Suppose that $V_{u_1} = A_\eta^*$; then $V_{u_1} b_1 (1 \otimes \Omega) = 1 \otimes \eta b_1$, hence $\epsilon(\vec{u})$ cancels unless $V_{u_2} \in \{V_0, A_\eta, P(T)\}$. Let $p = \min\{s : s > 1, u_s \neq 0\}$. It follows that $V_{u_p} \in \{A_\eta, P(T)\}$. Since the restrictions of $A_\eta, P(T)$ to $\mathcal{T}(\mathcal{H}) \otimes \mathcal{K}$ are 0, we have

$$\begin{aligned} V_{u_p} b_p \cdots V_{u_1} b_1 (1 \otimes \Omega) &= V_{u_p} b_p \left(V_0 b_{p-1} \cdots V_0 b_2\right) V_{u_1} b_1 (1 \otimes \Omega) \\ &= V_{u_p} b_p \varphi(X b_{p-1} \cdots X b_2) V_{u_1} b_1 (1 \otimes \Omega). \end{aligned}$$

If $V_{u_p} = P(T)$ and $s = \min\{q : p < q \leq n, u_q \neq 0\}$, from a similar argument as above we have that $u_{p+1} = \cdots = u_{q-1} = 0$ and

$$\begin{aligned} V_{u_q} b_q \cdots V_{u_1} b_1 (1 \otimes \Omega) \quad = \\ P(T) b_q \varphi(X b_{q-1} \cdots X b_{p+1}) P(T) b_p \varphi(X b_{p-1} \cdots X b_2) A_\eta^* b_1 (1 \otimes \Omega) \end{aligned}$$

Note also that, for all $b_1, \dots, b_m \in \mathcal{B}$, one has

$$\begin{aligned} T b_m \cdots T b_2 A_\eta^* b_1 (1 \otimes \Omega) &\in 1 \otimes \mathcal{K} \\ A_\eta b_m T b_{m-1} \cdots T b_2 A_\eta^* b_1 (1 \otimes \Omega) &\in 1 \otimes \Omega \mathcal{D}, \end{aligned}$$

hence $\epsilon(\vec{u})$ cancels unless there exists some $j > 1$ such that $V_{u_j} = A_\eta$.

Using the results above, we have that

$$\begin{aligned} \sum_{\substack{\vec{u} \in J(n) \\ u_1 = 4}} \epsilon(\vec{u}) \quad &= \quad \sum_{s=1}^{n} \sum_{\substack{p_1 \cdots < p_s \\ 1 < p_1, p_s \leq n}} \sum_{\substack{u \in J(n) \\ u_1 = 4, u_{p_s} = 2 \\ u_{p_l} = 1, l \neq s}} \epsilon(\vec{u}) \\ &= \sum_{s=1}^{n} \sum_{\substack{p_1 \cdots < p_s \\ 1 < p_1, p_s \leq n}} \theta(X b_n \cdots X b_{p_s + 1}) \cdot \theta(A_\eta b_{p_s} \varphi(X b_{p_s - 1} \cdots X b_{s-1} + 1) \cdot \\ &\quad P(T) b_{p_{s-1}} \varphi(X b_{p_{s-1} - 1} \cdots X b_{p_{s-2} + 1}) \cdots P(T) b_{p_1} \varphi(X b_{p_1} - 1 \cdots X b_2) A_\eta^* b_1) \end{aligned}$$

Comparing these relations with the definition of the c-free cumulants, we have q.e.d.. □

Theorem 4.8. *The pair* $(\mu, \nu) \in \Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$ *is* $\boxed{c}$*-infinitely divisible if and only if* ν *is* $\boxplus$*-infinitely divisible and the restriction of* ${}^c\rho_{\mu,\nu}$ *to* $\mathcal{B}\langle \mathcal{X} \rangle_0\rangle$ *satisfies property (1).*

Proof. Suppose that ν is infinitely divisible (hence, from Theorem 3.8, the restriction of ρ_ν to $\mathcal{B}\langle \mathcal{X} \rangle_0$ is positive) and that the restriction of ${}^c\rho_{\mu,\nu}$ to $\mathcal{B}\langle \mathcal{X} \rangle_0$ satisfies (1).

Let $\mathcal{H}$ be the left $\mathcal{B}$-bimodule $\mathcal{B}\langle \mathcal{X} \rangle_0$ with the pairing

$$\langle f(\mathcal{X}), g(\mathcal{X}) \rangle_{\mathcal{H}} = \rho_\nu (g(\mathcal{X})^* f(\mathcal{X}))$$

and $\mathcal{K}$ be the left $\mathcal{D}$-module $\mathcal{B}\langle \mathcal{X} \rangle_0 \otimes_{\mathcal{B}} \mathcal{D}$ with the pairing

$$\langle f(\mathcal{X}) \otimes d_1, g(\mathcal{X}) \otimes d_2 \rangle_{\mathcal{D}} = d_2^* \cdot {}^c\rho_{\mu,\nu}(g^*(\mathcal{X}) f(\mathcal{X})) d_1.$$

Since the maps ρ_ν and ${}^c\rho_{\mu,\nu}$ staify (1), we have that $\mathcal{H}$ and $\mathcal{K}$ are semi-inner-product $\mathcal{B}$-, respectively $\mathcal{D}$-modules.

Define

$$\begin{aligned} t \in \mathcal{L}(\mathcal{H}), &\qquad t(f(\mathcal{X})) = \mathcal{X} f(\mathcal{X}) \\ T \in \mathcal{L}(\mathcal{K}), &\qquad T(f(\mathcal{X}) \otimes d) = \mathcal{X} f(\mathcal{X}) \otimes d \end{aligned}$$

and note that t and T are selfadjoint maps. Also let $\lambda_1 = \rho_\nu(\mathcal{X}) \in \mathcal{B}$ and $\lambda_2 = {}^c\rho_{\mu.\nu}(\mathcal{X}) \in \mathcal{D}$.

As in Lemma 4.7, consider

$$\mathcal{E} = (\mathcal{T}(\mathcal{H}) \otimes_{\mathcal{B}} \mathcal{D}) \oplus (1_{\mathcal{B}} \otimes_{\mathcal{B}} \Omega\mathcal{D}) \oplus (\mathcal{T}(\mathcal{H}) \otimes_{\mathcal{B}} \mathcal{K})$$

and $V \in \mathcal{L}(\mathcal{E})$ given by

$$V = a_{\mathcal{X}} + a^*_{\mathcal{X}} + p(t) + \lambda_1 I_1 + A_{\mathcal{X}\otimes 1} + A^*_{\mathcal{X}\otimes 1} + \lambda_2 I_2.$$

Denoting $\varphi(\cdot) = \langle \cdot(1\otimes 1), 1\otimes 1\rangle$, $\theta(\cdot) = \langle \cdot(1\otimes\Omega), 1\otimes\Omega\rangle$ and applying Lemma 4.7 we have that the free and c-free cumulants of V with respect to (θ, φ) are given by:

$$\begin{aligned}
\kappa_{1,\mathcal{X}}(b_1) &= \lambda_1 b_1 = \rho_\nu(\mathcal{X}) \\
&= \kappa_{1,\nu}(b_1) \\
\kappa_{n,\mathcal{X}}(b_1, \dots, b_n) &= \langle a_{\mathcal{X}} b_1 p(T) b_2 \cdots p(t) b_{n-1} a^*_{\mathcal{X}} b_n (1\otimes 1), 1\otimes 1\rangle \\
&= \langle a_{\mathcal{X}} (b_1 \mathcal{X} b_2 \cdots \mathcal{X} b_{n-1} \mathcal{X} b_n \otimes 1), 1\otimes 1\rangle \\
&= \langle \rho_\nu(\mathcal{X} b_1 \cdots \mathcal{X} b_n) \otimes 1, 1\otimes 1\rangle \\
&= \kappa_{n,\nu}(b_1, \dots, b_n),
\end{aligned}$$

respectively by

$$\begin{aligned}
{}^c\kappa_{1,V}(b_1) &= \lambda_2 b_1 = {}^c\rho_{\mu,\nu}(\mathcal{X} b_1) \\
&= {}^c\kappa_{1,\mu,\nu}(b_1) \\
{}^c\kappa_{n,V}(b_1, \dots, b_n) &= \langle A_{\mathcal{X}\otimes 1} b_1 P(T) b_2 \cdots P(T) b_{n-1} A^*_{\mathcal{X}\otimes 1} b_1 (1\otimes\Omega), 1\otimes\Omega\rangle \\
&= \langle A_{\mathcal{X}\otimes 1}(b_1 \mathcal{X} b_2 \cdots \mathcal{X} b_n \otimes 1), 1\otimes\Omega\rangle \\
&= \langle {}^c\rho_{\mu,\nu}(\mathcal{X} b_1 \cdots \mathcal{X} b_n) \otimes \Omega, 1\otimes\Omega\rangle \\
&= {}^c\rho_{\mu,\nu}(\mathcal{X} b_1 \cdots \mathcal{X} b_n) \\
&= {}^c\kappa_{n,\mu,\nu}(b_1, \dots, b_n).
\end{aligned}$$

Fix $N > 0$ and define

$$V_N = \frac{1}{\sqrt{N}} a_{\mathcal{X}} + \frac{1}{\sqrt{N}} a^*_{\mathcal{X}} + p(t) + \frac{1}{N}\lambda_1 I_1 + \frac{1}{\sqrt{N}} A_{\mathcal{X}\otimes 1} + \frac{1}{\sqrt{N}} A^*_{\mathcal{X}\otimes 1} + P(T) + \frac{1}{N}\lambda_2 I_2.$$

Using again Lemma 4.7, similar computations as above give $(n \geq 1)$:

$$\begin{aligned}
\kappa_{n,V_N} &= \frac{1}{N}\kappa_{n,V} \\
{}^c\kappa_{n,V_N} &= \frac{1}{N}{}^c\kappa_{n,V}
\end{aligned}$$

hence q.e.d..

The converse implication is Proposition 4.6. □

5. The non-commutative $R-$ and ${}^cR$-transforms and non-commutative free Levy-Hincin formula

5.1. **The R- and ${}^cR-$transforms and free infinite divisibility: scalar case.**

Definition 5.1. *Let (μ,ν) be a pair of compactly supported measures on $\mathbb{R}$. If $M_\mu(z)$, $M_\nu(z)$ are the moment-generating series for μ, respectively ν, that is*

$$\begin{aligned} M_\mu(z) &= \sum_{n=0}^{\infty}\int_{\mathbb{R}} t^n d\mu(t) z^n \\ M_\nu(z) &= \sum_{n=0}^{\infty}\int_{\mathbb{R}} t^n d\nu(t) z^n \end{aligned}$$

then the R-transform of ν, respectively the ${}^cR$-transform of (μ,ν) are the analytic functions R_ν, ${}^cR_{\mu,\nu}$ given by

$$M_\nu(z) - 1 = R_\nu\left(zM_\nu(z)\right) \tag{10}$$

$$(M_\mu(z)-1)M_\nu(z) = M_\mu(z)\cdot {}^cR_{\mu,\nu}\left(zM_\nu(z)\right). \tag{11}$$

If A is a $*$-algebra and $\varphi : A \longrightarrow \mathbb{C}$ is a positive linear functional, then a selfadjoint element $a \in A$ determines a compactly supported real measure μ_a via

$$\int_{\mathbb{R}} t^n d\mu_a(t) = \varphi(a^n).$$

For convenience, we will also use the notation R_a for R_{μ_a}, respectively the notation ${}^cR_a$ for ${}^cR_{\mu_a,\nu_a}$ for the case that the the $*$-algebra A endowed with two positive linear functionals.

The key property of the R- and ${}^cR$-transforms is the linearization of the free convolution ([30], [22]), respectively c-free additive convolution: if a and b are free, respectively c-free, selfadjoint elements from A, then

$$R_{a+b}(z) = R_a(z) + R_b(z) \tag{12}$$

$${}^cR_{a+b} = {}^cR_a(z) + {}^cR_b(z) \tag{13}$$

The goal of this material is to give a non-commutative analogue for the following theorem ([22], Theorem 13.16 and [12], Theorem 5):

Theorem 5.2. *Let ν be a compactly supported supported probability measure on $\mathbb{R}$ and let $R_\nu(z) = \sum_{n=1}^{\infty}\kappa_n z^n$ be the Taylor expansion of its R-transform. Then the following statements are equivalent:*

1. *ν is infinitely divisible.*
2. *The sequence $\{\kappa_n\}_n \geq 2$ is positive definite, i.e. there exists some real measure σ such that*

   $$\kappa_n = \int_{\mathbb{R}} t^{n-2} d\sigma$$

3. *The R-transform of ν is of the form*

   $$\frac{1}{z}R_\nu(z) = \kappa_1 + \int_{\mathbb{R}} \frac{z}{1-tz} d\rho(t),$$

   for some finite measure ρ on $\mathbb{R}$ with compact support.

Moreover, if (μ,ν) is a pair of compactly supported measures on $\mathbb{R}$, then (μ,ν) is c-free infinitely divisible if and only if ν is infinitely divisible and ${}^cR_{\mu,\nu}$ satisfies the condition (3) from above.

5.2. **Non-Commutative functions.** For stating our main result, we introduce the language of noncommutative [11] or fully matricial [29] functions, see also the pioneering work [25, 26]. For a vector space $\mathcal{V}$ over $\mathbb{C}$, we let $\mathcal{V}^{n\times m} = \mathcal{V}\otimes_{\mathbb{C}} M_{n\times m}(\mathbb{C})$ denote $n \times m$ matrices over $\mathcal{V}$ (in literature - for example in [11] - on $\mathcal{V}^{n\times n}$ is used the algebra structure induced by the tensor algebra $\mathbf{T}(\mathcal{V})$ over $\mathcal{V}$; in order to avoid confusion, when $\mathcal{V}$ is an algebra, we will use the notation $M_n(\mathcal{V})$ for the *algebra* $M_n(\mathcal{V})$ of $n \times n$ matrices over $\mathcal{V}$). We define the *noncommutative space* over $\mathcal{V}$ by $\mathcal{V}_{\mathrm{nc}} = \coprod_{n=1}^{\infty} \mathcal{V}^{n\times n}$. We call $\Omega \subseteq \mathcal{V}_{\mathrm{nc}}$ a *noncommutative set* if it is closed under direct sums. Explicitly, denoting $\Omega_n = \Omega \cap \mathcal{V}^{n\times n}$, we have

$$a \oplus b = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \in \Omega_{n+m}$$

for all $a \in \Omega_n$, $b \in \Omega_m$. Notice that matrices over $\mathbb{C}$ act from the right and from the left on matrices over $\mathcal{V}$ by the standard rules of matrix multiplication.

A noncommutative set $\Omega \subseteq \mathcal{V}_{\mathrm{nc}}$ is called *upper admissible* if for all $a \in \Omega_n$, $b \in \Omega_m$ and all $c \in \mathcal{V}^{n\times m}$, there exists $\lambda \in \mathbb{C}$, $\lambda \neq 0$, such that

$$\begin{bmatrix} a & \lambda c \\ 0 & b \end{bmatrix} \in \Omega_{n+m}.$$

This notion is crucial since it is used to define the (right) noncommutative difference-differential operators by evaluating a noncommutative function on block upper triangular matrices. We will encounter only the following upper admissible noncommutative sets:

1. The set $\mathrm{Nilp}(\mathcal{V}) = \coprod_{n=1}^{\infty} \mathrm{Nilp}(\mathcal{V}; n)$ of nilpotent matrices over $\mathcal{V}$. Here the set $\mathrm{Nilp}(\mathcal{V}; n)$ of nilpotent $n\times n$ matrices over $\mathcal{V}$ consists of all $a \in \mathcal{V}^{n\times n}$ such that $a^r = 0$ for some r, where we view a as a matrix over the tensor algebra $\mathbf{T}(\mathcal{V})$ of $\mathcal{V}$ over $\mathbb{C}$. This is equivalent to tat^{-1} being strictly upper triangular for some $t \in M_n(\mathbb{C})$ (the equivalence follows from Engel's Theorem — notice that we can restrict ourselves to the finite dimensional subspace of $\mathcal{V}$ spanned by the elements of a).
2. A noncommutative ball $\mathbb{B}(\mathcal{A}, \rho) = \{a \in \mathcal{A}_{\mathrm{nc}} \colon \|a\| < \rho\}$ centered in zero and of radius $\rho > 0$ over a C*-algebra $\mathcal{A}$ ($\mathcal{A}$ could have been replaced by any operator space with the corresponding operator space norm).
3. The upper and lower fully matricial half-planes $\mathbb{H}^+(A_{\mathrm{nc}})$ and $\mathbb{H}^-(A_{\mathrm{nc}})$ over a C*-algebra A, where if $\mathcal{C}$ is a C*-algebra, then

$$\begin{aligned} \mathbb{H}^+(\mathcal{C}) &= \{a \in \mathcal{C}, \Im a = \frac{(a-a^*)}{2} > 0\} \\ \mathbb{H}^-(\mathcal{C}) &= \{a \in \mathcal{C}, \Im a = \frac{(a-a^*)}{2} < 0\} \end{aligned}$$

   and $\mathbb{H}^{\pm}(A_{\mathrm{nc}}) = \coprod_{n=1}^{\infty} \mathbb{H}^{\pm}(M_n(A))$.

Let $\mathcal{V}$ and $\mathcal{W}$ be vector spaces over $\mathbb{C}$, and let $\Omega \subseteq \mathcal{V}_{\mathrm{nc}}$ be a noncommutative set. A mapping $f\colon \Omega \to \mathcal{W}_{\mathrm{nc}}$ with $f(\Omega_n) \subseteq \mathcal{W}^{n\times n}$ is called a *noncommutative function* if f satisfies the following two conditions:

- f *respects direct sums*: $f(a \oplus b) = f(a) \oplus f(b)$ for all $a, b \in \Omega$.
- f *respects similarities*: if $a \in \Omega_n$ and $s \in \mathbb{C}^{n\times n}$ is invertible with $sas^{-1} \in \Omega_n$, then $f(sas^{-1}) = sf(a)s^{-1}$.

We will denote $f_n = f|_{\Omega_n} \colon \Omega_n \to \mathcal{W}^{n\times n}$. While we will not need this fact, it is important to notice that the two conditions in the definition of a noncommutative function can be actually replaced by a single one: a mapping $f\colon \Omega \to \mathcal{W}_{\mathrm{nc}}$ with $f(\Omega_n) \subseteq \mathcal{W}^{n\times n}$ respects direct sums and similarities if and only if it *respects intertwinings*: for any $a \in \Omega_n$, $b \in \Omega_m$, and $t \in \mathbb{C}^{n\times m}$ such that $at = tb$, one has $f(a)t = tf(b)$. This last condition goes back to [25, 26].

Let $\alpha\colon \mathcal{V}^k \longrightarrow \mathcal{W}$ be a multilinear mapping; we set

$$\widetilde{\alpha} = \widetilde{\alpha}_n \overset{\mathrm{def}}{=} \alpha \otimes \mathrm{Id}_n \colon \left(\mathcal{V}^{\otimes k}\right)^{n\times n} \longrightarrow \mathcal{W}^{n\times n}$$

for each $n \in \mathbb{N}$. It is clear that the mapping $a \mapsto \widetilde{\alpha}(a^k)$, where we view a as a matrix over $\mathbf{T}(\mathcal{V})$, respects direct sums and similarities, so that it defines a noncommutative function from $\mathcal{V}_{\mathrm{nc}}$ to $\mathcal{W}_{\mathrm{nc}}$. Therefore for any linear mapping $\phi\colon \mathbf{T}(\mathcal{V}) \longrightarrow \mathcal{W}$, we obtain a noncommutative function defined by

$$f(a) = \widetilde{\phi}\left((\mathbb{1} - a)^{-1}\right) = \sum_{k=0}^{\infty} \widetilde{\phi}(a^k) \tag{14}$$

(where the notation $\mathbb{1}$ should be understood as Id_n componentwise, i.e. $f(a) = \widetilde{\phi}_n\left((\mathrm{Id}_n - a)^{-1}\right)$ for $a \in \mathcal{V}^{n\times n}$), except that we have to make sense of the infinite sum on the right-hand side.

1. If $a \in \mathrm{Nilp}(\mathcal{V})$ then the sum is finite, so that f is always a noncommutative function on $\mathrm{Nilp}(\mathcal{V})$.
2. If $\phi\colon \mathbf{T}(\mathcal{A}) \longrightarrow \mathcal{C}$, where $\mathcal{A}$ and $\mathcal{C}$ are C^* algebras, and we have an exponential growth estimate: $\|\phi|_{\mathcal{A}^{\otimes k}\to\mathcal{C}}\|_{\mathrm{cb}} \leq \alpha\beta^k$ (where $\mathcal{A}^{\otimes k}$ is considered with the Haagerup tensor norm (see [16], Chapter 17) and $\|\cdot\|_{\mathrm{cb}}$ denotes the completely bounded norm), then the series defining f converges absolutely and uniformly on any noncommutative ball $\mathbb{B}(\mathcal{A}, r)$ over $\mathcal{A}$ of radius $r < 1/\beta$, so that f is a noncommutative function on the noncommutative ball $\mathbb{B}(\mathcal{A}, 1/\beta)$.

There is — in a sense — a converse to this construction that we briefly describe, though we will make no real use of it here. A noncommutative function f admits a series expansion

$$f(a) = \sum_{k=0}^{\infty} \Delta_R^k f\widetilde{(\underbrace{0,\ldots,0}_{k+1})}(a^{\otimes k}). \tag{15}$$

More precisely:

1. If f is a noncommutative function on $\mathrm{Nilp}(\mathcal{V})$, then the sum is finite and the equality holds everywhere.
2. If f is a noncommutative function on a noncommutative ball $\mathbb{B}(\mathcal{A}, \rho)$ over a C^* algebra $\mathcal{A}$ with values in a noncommutative space $\mathcal{C}_{\mathrm{nc}}$ over a C^* algebra $\mathcal{C}$, which is bounded on noncommutative balls of radius less then ρ, then the series converges to f, uniformly on every noncommutative ball of radius less than ρ. (The convergence still holds if f is only assumed to be locally bounded in every matrix dimension separately, but then it is no longer uniform across matrix dimensions.)

The multilinear forms $\Delta_R^k f(\underbrace{0,\ldots,0}_{k+1})\colon \mathcal{V}^k \longrightarrow \mathcal{W}$ are the values at $(0,\ldots,0)$ of the kth order noncommutative difference-differential operators applied to f. They are uniquely determined, and can be calculated directly by evaluating f on upper triangular matrices:

$$f\left(\begin{bmatrix} 0 & a_1 & 0 & \dots & 0 \\ 0 & 0 & a_2 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & a_k \\ 0 & 0 & 0 & \dots & 0 \end{bmatrix}\right)$$

$$= \begin{bmatrix} f(0) & \Delta_R f(0,0)(a_1) & \cdots & \cdots & \Delta_R^k f(0,\dots,0)(a_0,\dots,a_k) \\ 0 & f(0) & \ddots & & \Delta_R^{k-1} f(0,\dots,0)(a_2,\dots,a_k) \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & f(0) & \Delta_R f(0,0)(a_k) \\ 0 & \cdots & \cdots & 0 & f(0) \end{bmatrix}.$$

5.3. The non-commutative B-, R- and ${}^cR$-transforms.

In particular, we can apply the construction (14) to $\mu \in \Sigma_{\mathcal{B}:\mathcal{D}}$, and define

$$M_\mu(b) = \widetilde{\mu}\left((\mathbb{1} - \mathcal{X}b)^{-1}\right) = \sum_{k=0}^{\infty} \widetilde{\mu}\left((\mathcal{X}b)^n\right),$$

where the notation $\mathbb{1}$ is the one from the previous section and, as before, we identify the monomials $(\mathcal{X}b)^n$ from $\mathcal{B}\langle\mathcal{X}\rangle_0$ to $b^{\otimes n}$ from the tensor algebra $\mathbf{T}(\mathcal{B})$. M_μ is a noncommutative function with values in $\mathcal{D}_{\mathrm{nc}}$. It is always defined on $\mathrm{Nilp}(\mathcal{B})$. If $\mathcal{D}$ is a C*-algebra and $\mu \in \Sigma^0_{\mathcal{B}:\mathcal{D}}$, that is there is some $C > 0$ such that $\|\mu_k\|_{\mathrm{cb}} \leq C^{k+1}$, where μ_k is the restriction of μ to the subspace of $\mathcal{B}\langle\mathcal{X}\rangle$ spanned by words with exactly k occurrences of the symbol X (that can be identified with $\mathcal{B}^{\otimes k}$), then M_μ is also defined on a noncommutative ball in $\mathcal{B}$ with sufficiently small radius. Remark that (14) also implies $\Delta_R^k M_\mu(0,\dots,0) = \mu_k$.

Consider now $\nu \in \Sigma_{\mathcal{B}}$ and $\mu \in \Sigma_{\mathcal{B}:\mathcal{D}}$. We define the linear maps ρ_ν, ${}^c\rho_{\mu,\nu}$ as in Section 3, respectively Section 4, that is the $\mathcal{B}$-bimodule extensions of

$$\rho_\nu(\mathcal{X}b_1 \cdots \mathcal{X}b_p) = \kappa_{\nu,p}(b_1,\dots,b_p), \qquad {}^c\rho_{\mu,\nu}(\mathcal{X}b_1 \cdots \mathcal{X}b_p) = {}^c\kappa_{\mu,\nu,p}(b_1,\dots,b_p)$$

and the map β_μ as the $\mathcal{B}$-bimodule extension of

$$\beta_\mu(\mathcal{X}b_1 \cdots \mathcal{X}b_p) = B_{\mu,p}(b_1,\cdot,b_p).$$

Using again (14), we define the noncommutative functions R_ν (with values in $\mathcal{B}_{\mathrm{nc}}$) and B_ν, ${}^cR_{\mu,\nu}$ (with values in $\mathcal{D}_{\mathrm{nc}}$) via the equations

$$\begin{aligned} R_\nu(b) &= \widetilde{\rho_\nu}\left((\mathbb{1} - \mathcal{X}b)^{-1}\right) - \mathbb{1} \\ {}^cR_{\mu,\nu}(b) &= \widetilde{{}^c\rho_{\mu,\nu}}\left((\mathbb{1} - \mathcal{X}b)^{-1}\right) - \mathbb{1} \\ B_\mu(b) &= \widetilde{\beta_\mu}\left((\mathbb{1} - \mathcal{X}b)^{-1}\right) - \mathbb{1}. \end{aligned}$$

Remark again that (14) implies $\Delta_R^p \mathcal{R}_\nu(0,\dots,0) = \kappa_{\nu,p}$, $\Delta_R^p {}^c\mathcal{R}_{\mu,\nu}(0,\dots,0) = {}^c\kappa_{\mu,\nu,p}$ and $\Delta_R^p B_\mu(0,\dots,0) = B_{\mu,p}$.

Proposition 5.3. *For all* $(\mu, \nu) \in \Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$ *we have that*

$$\begin{aligned} Id_n \otimes \rho_\nu &= \rho_{\nu^{(n)}} \\ Id_n \otimes {}^c\rho_{\mu,\nu} &= \rho_{\mu^{(n)},\nu^{(n)}} \\ Id_n \otimes \beta_\mu &= \beta_{\mu^{(n)}}. \end{aligned}$$

Proof. For the first property, it suffices to show that for all positive integers m, all $1 \leq i_k, j_k \leq n$ $(k = 1, \dots, n)$ and all $b_1, \dots, b_m \in \mathcal{B}$ we have that

$$\mathrm{Id}_n \otimes \rho_\nu(\mathcal{X}B_1 \cdots \mathcal{X}B_m) = \rho_{\nu^{(n)}}(\mathcal{X}B_1 \cdots \mathcal{X}B_m),$$

where $B_k = e_{i_k,j_k} \otimes b_k$, for e_{i_k,j_k} the complex $n \times n$ matrix with the (i_k, j_k) entry 1 and all others 0.

For $m = 1$ the property is trivial. Suppose now that the assertion holds true for all $m < N$. The definition of free cumulants gives

$$\begin{aligned} \rho_{\nu^{(n)}}(\mathcal{X}B_1 \cdots \mathcal{X}B_N) &= \nu^{(n)}(\mathcal{X}B_1 \cdots \mathcal{X}B_N) - \\ \sum_{p=1}^{N-1} \sum_{\substack{s_1 < \cdots < s_p \\ s_1 = 1, s_p \leq N}} & \rho_{\nu^{(n)}}\Big(\mathcal{X}B_1 \nu^{(n)}(\mathcal{X}B_2 \cdots \mathcal{X}B_{s_2-1}) \cdots \mathcal{X}B_{s_p} \nu^{(n)}(\mathcal{X}B_{s_p+1} \cdots \mathcal{X}B_N)\Big). \end{aligned}$$

From the induction hypothesis, the right-hand side cancels unless $i_k = J_{k-1}$ for all $1 < k \leq N$; in this case, the equation above becomes

$$\begin{aligned} \rho_{\nu^{(n)}}(\mathcal{X}B_1 \cdots \mathcal{X}B_N) &= e_{i_1,j_N} \otimes \nu(\mathcal{X}b_1 \cdots \mathcal{X}b_N) - \\ \sum_{p=1}^{N-1} \sum_{\substack{s_1 < \cdots < s_p \\ s_1 = 1, s_p \leq N}} & \rho_{\nu^{(n)}}\left(e_{i_1,j_N} \otimes [\mathcal{X}b_1 \nu(\mathcal{X}b_2 \cdots \mathcal{X}b_{s_2-1}) \cdots \mathcal{X}b_{s_p} \nu(\mathcal{X}b_{s_p+1} \cdots \mathcal{X}b_N)]\right) \\ &= e_{i_1,j_N} \otimes \rho_\nu(\mathcal{X}b_1 \cdots \mathcal{X}b_N) \\ &= \mathrm{Id}_n \otimes \rho_\nu(\mathcal{X}B_1 \cdots \mathcal{X}B_N) \end{aligned}$$

hence the conclusion.
The argument for ${}^c\rho_{\mu,\nu}$ and β_μ is analogous. ☐

Proposition 5.3 implies that

$$R_\nu^{(n)}(b) = \sum_{n=1}^{\infty} \kappa_{n,\nu^{(n)}}(b, \dots, b)$$

and the analogous relations for the components of ${}^cR_{\mu,\nu}$ and B_μ. From the moment-cumulant recursions, we have then the following

Corollary 5.4. *The non-commutative functions* R_ν, B_ν *and* ${}^cR_{\mu,\nu}$ *satisfy the equations:*

$$M_\mu(b) - \mathbb{1} = B_\mu(b) \cdot M_\mu(b) \tag{16}$$

$$M_\nu(b) - \mathbb{1} = R_\nu\left(bM_\nu(b)\right) \tag{17}$$

$$(M_\mu(b) - \mathbb{1}) \cdot M_\nu(b) = M_\mu(b) \cdot {}^cR_{\mu,\nu}\left(bM_\nu(b)\right). \tag{18}$$

Remark 5.5. *If* $(\mu, \nu) \in \Sigma^0_{\mathcal{B}:\mathcal{D}} \times \Sigma^0_{\mathcal{B}}$ *then* R_ν, ${}^cR_{\mu,\nu}$ *and* B_μ *are well-defined on a non-commutative ball (as described in Section 5.2) centered in* 0.

Proof. For B_μ, the assertion is trivial since $B_\mu(b)$ is well-defined if M_μ is invertible. For R_ν and ${}^cR_{\mu,\nu}$ we will use combinatorial techniques, namely the Moebius inversion formula for the partially ordered set of non-crossing partitions. First, we need the following general property (the Moebius inversion formula).

Proposition: *Let P be a finite partially ordered set and K a complex vector space. Then there exist a map* $\operatorname{moeb} : P \times P \longrightarrow \mathbb{R}$ *such that if the maps $f, g : P \longrightarrow K$ have the property*

$$f(\pi) = \sum_{\substack{\sigma \in P \\ \sigma \leq \pi}} g(\sigma), \pi \in P$$

then

$$g(\pi) = \sum_{\substack{\sigma \in P \\ \sigma \leq \pi}} f(\sigma) \cdot \operatorname{moeb}(\sigma, \pi).$$

We will apply the above result on the partially ordered set $\mathcal{NC}$ of non-crossing partitions. By a partition on the ordered set $\langle n \rangle = \{1, 2, \ldots, n\}$ we will understand a collection of mutually disjoint subsets of $\langle n \rangle$, $\gamma = (B_1, \ldots, B_q)$ called *blocks* whose union is the entire set $\langle n \rangle$. A crossing is a sequence $i < j < k < l$ from $\langle n \rangle$ with the property that there exist two different blocks B_r and B_s such that $i, k \in B_r$ and $j, l \in B_s$. A partition that has no crossings will be called non-crossing. The set of all non-crossing partitions on $\langle n \rangle$ will be denoted by NC(n). For $\gamma \in NC(n)$ a block $B = (i_1, ..., i_k)$ of γ will be called *interior* if there exists another block $D \in \gamma$ and $i, j \in D$ such that $i < i_1, i_2, \ldots, i_k < j$. A block will be called *exterior* if is not interior. Each $NC(n)$ has a lattice structure with respect to block refinement with biggest element 1_m- the partition with a single block; the corespondent Moebius function satisfies $|\operatorname{moeb}(\pi, 1_n)| \leq 4^n$ (see [22], Lecture 13). Define $\mathcal{NC} = \coprod_{n=1}^{\infty} NC(n)$.

Finally, let $(\mu, \nu) \in \Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$, $b \in \mathcal{B}$. We define $f, F : \mathcal{NC} \longrightarrow \mathcal{D}$ as follows:

(a) $f(\pi_1 \cdot \pi_2) = f(\pi_1) \cdot f(\pi_2)$ and $F(\pi_1 \cdot \pi_2) = F(\pi_1) \cdot F(\pi_2)$,
where $\pi_1 \cdot \pi_2 \in NC(m+n)$ is obtained by juxtaposing $\pi_1 \in NC(n)$ and $\pi_2 \in NC(m)$.

(b) $f(1_m) = \nu((\mathcal{X}b)^m)$ and $F(1_m) = \mu((\mathcal{X}b)^m)$,
where $1_m \in NC(m)$ is the partition with a single block $(1, 2, \ldots, m)$.

(c) $f(\overline{|\pi_1|\pi_2|\ldots|\pi_q|}) = \nu(\mathcal{X}b \cdot f(\pi_1) \cdot \mathcal{X}b \cdot f(\pi_2) \cdots \mathcal{X}b \cdot f(\pi_q) \cdot \mathcal{X}b)$
$F(\overline{|\pi_1|\pi_2|\ldots|\pi_q|}) = \mu(\mathcal{X}b \cdot f(\pi_1) \cdot \mathcal{X}b \cdot f(\pi_2) \cdots \mathcal{X}b \cdot f(\pi_q) \cdot \mathcal{X}b)$
where $\overline{|\pi_1|\pi_2|\ldots|\pi_q|}$ is the partition with a single exterior block with $q+1$ elements and with restrictions between the elements of the exterior block $\pi_1, \ldots, \pi_q$.

Similarly, we define $g, G : NC \longrightarrow \mathcal{D}$ as above, replacing μ with ${}^c\rho_{\mu,\nu}$ and ν with ρ_ν. With this notations, the moment-cumulant relation from Definition 3.4 amounts to

$$F(\pi) = \sum_{\substack{\sigma \in \mathcal{NC} \\ \sigma \leq \pi}} G(\sigma) \tag{19}$$

and applying the Moebius function property to equation 19, we get

$$G(\pi) = \sum_{\substack{\sigma \in \mathcal{NC} \\ \sigma \leq \pi}} F(\sigma) \cdot \operatorname{moeb}(\sigma, \pi).$$

Suppose now that $(\mu, \nu) \in \Sigma^0_{\mathcal{B}:\mathcal{D}} \times \Sigma^0_{\mathcal{B}}$ and that that $||\mu_n||_{cb}, ||\nu_n||_{cb} < M^{n+1}$ for all $n > 0$, henceforth $||F(\sigma)|| \leq M^{m+1}||b||^m$ for all $\sigma \in NC(m)$. Since ${}^c\kappa_{\mu,\nu,m}(b) = G(1_m)$, we have

$$\begin{aligned}
\|{}^c\kappa_{\mu,\nu,m}(b)\| &\leq \sum_{\sigma \in NC(m)} \|F(\sigma)\| \cdot |\mathrm{moeb}(\sigma, 1_m)| \\
&< (\sharp NC(m)) \cdot M^{m+1}\|b\|^m \cdot 4^m \\
&< \cdot M^{m+1}\|b\|^m \cdot 16^m, \ \text{since} \sharp NC(m) < 4^m.
\end{aligned}$$

Finally, $\kappa_{\nu,m}(b) = {}^c\kappa_{\mu,\nu,m}(b)$, so the assertion is proven also for the R-transform. □

5.4. **Main results.** The following property of the B-transform is a non-commutative analogue of Proposition 3.1. from [23], (i. e. the Nevalinna-Pick representation for the self-energy function of a real measure):

Theorem 5.6. *Let $\mathcal{B}$ be a C^*-algebra, $\mathcal{B} \subset \mathcal{D}$ be a unital inclusion of C^*-algebras and $\mu \in \Sigma_{\mathcal{B}:\mathcal{D}}$. Then there exists a selfadjoint $\alpha \in \mathcal{D}$ and a $\mathbb{C}$-linear map $\sigma : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}$, satisfying property (1) such that*

$$B_\mu(b) = \left[\alpha \cdot \mathbb{1} + \widetilde{\sigma}\left(b(\mathbb{1} - \mathcal{X}b)^{-1}\right)\right] \cdot b.$$

Moreover, if $\mu \in \Sigma^0_{\mathcal{B}:\mathcal{D}}$, then the moments of ν do not grow faster than exponentially (i.e. ν satisfies property (2).

Proof. Define the map $\beta_\mu : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}$ as the $\mathcal{B}$-bimodule extension of

$$\beta_\mu(\mathcal{X}b_1\mathcal{X} \cdots \mathcal{X}b_n) = B_{n,\mu}(b_1, \dots, b_n).$$

Let $\mathcal{K} = \mathcal{B}\langle\mathcal{X}\rangle \otimes_{\mathcal{B}} \mathcal{D}$. As shown in the proof of Theorem 2.5, equation (5), there exists some $\xi \in \mathcal{K}$ and a selfadjoint map $T \in \mathcal{L}(\mathcal{K})$ such that

$$B_{n,\mu}(b_1, \dots, b_n) = \langle b_1Tb_2 \cdots Tb_{n-1}(\xi b_n), \xi\rangle$$

If $b_n = 1$, the above relation becomes

$$\begin{aligned}
\langle b_1Tb_2 \cdots Tb_{n-1}\xi, \xi\rangle &= B_{n,\mu}(b_1, \dots, b_{n-1}, 1) \\
&= \beta_\mu(\mathcal{X}b_1\mathcal{X} \cdots \mathcal{X}b_{n-1}\mathcal{X})
\end{aligned}$$

Hence for all $f(\mathcal{X}) \in \mathcal{B}\langle\mathcal{X}\rangle$ we have that $\beta_\mu(\mathcal{X}f(\mathcal{X})\mathcal{X}) = \langle f(T)\xi, \xi\rangle$. It follows that the map $\sigma : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}$ given by $\sigma(f(\mathcal{X})) = \beta_\mu(\mathcal{X}f(\mathcal{X})\mathcal{X})$ satisfies property (1) and

$$B_{n,\mu}(b, \dots, b) = \sigma\left([\mathcal{X}b]^{n-1}\right) \cdot b,$$

that is the conclusion for $\alpha = B_{1,\mu}$. The last part is a trivial consequence of the fact that $\mu \in \Sigma^0_{\mathcal{B}:\mathcal{D}}$ implies that $\beta_\mu \in \Sigma^0_{\mathcal{B}:\mathcal{D}}$. □

Remark 5.7. If $\mu \in \Sigma^0_{\mathcal{B}:\mathcal{D}}$, the above result can be more explicit formulated in terms of the generalized rezolvent of μ from [28] and [2]. More precisely, for $\mu \in \Sigma^0_{\mathcal{B}:\mathcal{D}}$, its *generalized rezolvent* or *operator-valued Cauchy transform* is defined via

$$\mathcal{G}_\mu(b) = \widetilde{\mu}\big([b - \mathbb{1} \cdot \mathcal{X}]^{-1}\big).$$

As shown in [28] and [2], $\mathcal{G}_\mu$ is a non-commutative function, well-defined on $\mathbb{H}^+(\mathcal{B}_{\mathrm{nc}})$ and $\mathcal{G}_\mu(\mathbb{H}^+(M_n(\mathcal{B}))) \subseteq \mathbb{H}^-(M_n(\mathcal{D}))$. Hence, its reciprocal, $b \mapsto \left[\mathcal{G}_\mu(b)\right]^{-1}$ is also a non-commutative function, well defined on $\mathbb{H}^+(\mathcal{B}_{\text{nc}})$. Moreover, identifying the coefficients, we have that

$$\mathbb{1} - \big[\mathcal{G}_\mu(b)\big]^{-1} \cdot b^{-1} = B_\mu(b^{-1}) \tag{20}$$

for $b \in \mathbb{H}^+(B_{\text{nc}})$ with $||b^{-1}|| < M$ for some $M > 0$.

Using equation 20 and Theorem 5.6, we obtain that, for b as above,

$$\begin{aligned} b - \big[\mathcal{G}_\mu(b)\big]^{-1} &= \alpha \cdot \mathbb{1} + \widetilde{\sigma}(b^{-1}[\mathbb{1} - \mathcal{X} \cdot b^{-1}]^{-1}) \\ &= \alpha \cdot \mathbb{1} + \widetilde{\sigma}([(\mathbb{1} - \mathcal{X} \cdot b^{-1}) \cdot b]^{-1}) \\ &= \alpha \cdot \mathbb{1} + \widetilde{\sigma}([b - \mathbb{1} \cdot \mathcal{X}]^{-1}), \end{aligned}$$

The map $b \mapsto \widetilde{\sigma}([b - \mathbb{1} \cdot \mathcal{X}]^{-1})$ extends to $\mathbb{H}^+(\mathcal{B}_{\text{nc}})$ (see again [28] and [29]) therefore we obtained that the *operator-valued selfenergy function* of μ ($b \mapsto b - \big[\mathcal{G}_\mu(b)\big]^{-1}$) is the translate with a selfadjoint of the operator-valued Cauchy transform of some $\mathbb{C}$-linear map from $\mathcal{B}\langle\mathcal{X}\rangle$ to $\mathcal{D}$ satisfying properties 1 and 2 □

For the main result of this section, Theorem 5.10, we will first need the following lemma:

Lemma 5.8. *Let $\mathcal{B} \subset \mathcal{D}$ be a unital inclusion of C*-algebras and $\rho : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}$ be a unital $\mathcal{B}$-bimodule map. Then the restriction of ρ to $\mathcal{B}\langle\mathcal{X}\rangle_0$ satisfies property (1) if and only if there exists some ($\mathbb{C}$)-linear map $\sigma = \sigma(\rho) : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}$ satisfying property (1) such that $\sigma(f(\mathcal{X})) = \rho(\mathcal{X} f(\mathcal{X})\mathcal{X})$ for all $f(\mathcal{X}) \in \mathcal{B}\langle\mathcal{X}\rangle$.*

Proof. Suppose that first that the restriction of $\rho_{|\mathcal{B}\langle\mathcal{X}\rangle_0}$ satisfies (1) and define σ via $\sigma(f(\mathcal{X})) = \rho(\mathcal{X} f(\mathcal{X})\mathcal{X})$. If $\{f_j(\mathcal{X})\}_{j=1}^n$ is some family from $\mathcal{B}\langle\mathcal{X}\rangle_0$, then

$$\begin{aligned} \big[\sigma(f_j(\mathcal{X})^* f_i(\mathcal{X}))\big]_{i,j=1}^n &= \big[\rho(\mathcal{X} f_j(\mathcal{X})^* f_i(\mathcal{X})\mathcal{X})\big]_{i,j=1}^n \\ &= \big[\rho\big([f_j(\mathcal{X})\mathcal{X}]^*[f_i(\mathcal{X})\mathcal{X}]\big)\big]_{i,j=1}^n \geq 0 \end{aligned}$$

For the converse, note that σ satisfies (1) implies (cf [16], pag. 42) that $\mathcal{B}\langle\mathcal{X}\rangle \otimes \mathcal{D}$ is a semi-inner product $\mathcal{D}$-module with respect to the pairing generated by

$$\langle f(\mathcal{X}) \otimes d_1, g(\mathcal{X}) \otimes d_2 \rangle = d_2^* \sigma(g(\mathcal{X})^* f(\mathcal{X})) d_1.$$

Fix now a family $\{f_j(\mathcal{X})\}_{j=1}^n$ from $\mathcal{B}\langle\mathcal{X}\rangle_0$. Each $f_j(\mathcal{X})$ can be written as

$$f_j(\mathcal{X}) = \sum_{k=1}^{N(j)} g_{k,j}(\mathcal{X}) \cdot \mathcal{X} \cdot \alpha_{k,j}$$

with $g_{k,j}(\mathcal{X}) \in \mathcal{B}\langle\mathcal{X}\rangle$ and $\alpha_{k,j} \in \mathcal{B}$. Denote $\eta_j = \sum_{k=1}^{N(j)} g_{k,j}(\mathcal{X})\mathcal{X}\alpha_{k,j} \in \mathcal{B}\langle\mathcal{X}\rangle \otimes \mathcal{D}$. Then

$$\begin{aligned} \big[\rho(f_j(\mathcal{X})^* f_i(\mathcal{X}))\big]_{i,j=1}^n &= \big[\sum_{k=1}^{N(j)} \sum_{l=1}^{N(i)} \rho(\alpha_{k,j}^* \cdot \mathcal{X} \cdot g_{k,j}(\mathcal{X})^* g_{l,i}(\mathcal{X}) \cdot \mathcal{X} \cdot \alpha_{l,i})\big]_{i,j=1}^n \\ &= \big[\sum_{k=1}^{N(j)} \sum_{l=1}^{N(i)} \alpha_{k,j}^* \cdot \sigma(g_{k,j}(\mathcal{X})^* g_{l,i}(\mathcal{X})) \cdot \alpha_{l,i}\big]_{i,j=1}^n \\ &= \big[\langle \sum_{l=1}^{N(i)} g_{l,i}(\mathcal{X}) \otimes \alpha_{l,i}, \sum_{k=1}^{N(j)} g_{k,j}(\mathcal{X}) \otimes \alpha_{k,j} \rangle\big]_{i,j=1}^n \\ &= \big[\langle \eta_i, \eta_j \rangle\big]_{i,j=1}^n \geq 0. \end{aligned}$$

□

An immediate consequence of Theorem 5.6 and the above Lemma is the following

Corollary 5.9. *If $\mu \in \Sigma_{\mathcal{B}:\mathcal{D}}$ then $\beta_{\mu|\mathcal{B}\langle\mathcal{X}\rangle_0}$ satisfies property* (1).

Theorem 5.10. *As before, $\mathcal{B} \subset \mathcal{D}$ will be a unital inclusion of unital C$*$-algebras.*

(i)*Let $\mu \in \Sigma_{\mathcal{B}}$. Then μ is $\boxplus$-infinitly divisible if and only if there exist some selfadjoint $\alpha \in \mathcal{B}$ and some $\mathbb{C}$-linear map $\sigma : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{B}$ satisfying property* (1), *such that*

$$R_\mu(b) = \left[\alpha \cdot \mathbb{1} + \widetilde{\sigma}\left(b(\mathbb{1} - \mathcal{X}b)^{-1}\right)\right] \cdot b.$$

(ii) *Let $(\mu, \nu) \in \Sigma_{\mathcal{B}:\mathcal{D}} \times \Sigma_{\mathcal{B}}$. Then (μ, ν) is $\boxed{c}$-infinitely divisible if and only if ν is $\boxplus$-infinitely divisible and there exist some selfadjoint $\alpha \in \mathcal{B}$ and some $\mathbb{C}$-linear map $\sigma : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}$ satisfying property* (1) *such that*

$${}^cR_{\mu,\nu}(b) = \left[\alpha \cdot \mathbb{1} + \widetilde{\sigma}\left(b(\mathbb{1} - \mathcal{X}b)^{-1}\right)\right] \cdot b.$$

Moreover, if $\mu \in \Sigma^0_{\mathcal{B}}$, respectively $(\mu, \nu) \in \Sigma^0_{\mathcal{B}:\mathcal{D}}$, then the moments of the correspondent maps σ do not grow faster than exponentially.

Proof. Since $\Sigma_{\mathcal{B}:\mathcal{B}} = \Sigma_{\mathcal{B}:\mathcal{D}}$, it suffices to prove the assertions for ${}^cR_{\mu,\nu}$.

Since ${}^c\rho_{\mu,nu}(\mathcal{X}b_1 \cdots \mathcal{X}b_n) = \kappa_{\mu,n}(b_1, \ldots, b_n)$, Proposition 5.3 implies that

$${}^cR_{\mu,\nu}(b) = M_{{}^c\rho_{\mu,\nu}}(b) - \mathbb{1}.$$

for all $b \in Nilp(\mathcal{B})$ or for b or sufficiently small norm if $(\mu, \nu) \in \Sigma^0_{\mathcal{B}:\mathcal{D}} \times \Sigma^0_{\mathcal{B}}$.

Identifying $\mathcal{X}$ to $\mathbb{1} \cdot \mathcal{X}$, we have that

$$\begin{aligned}
{}^cR_{\mu,\nu}(b) &= \sum_{n=1}^{\infty} \widetilde{{}^c\rho_{\mu,\nu}}\left((\mathcal{X}b)^n\right) \\
&= \widetilde{{}^c\rho_{\mu,\nu}}(\mathcal{X}) \cdot b + \left[\sum_{n=2}^{\infty} \widetilde{{}^c\rho_{\mu,\nu}}\left((\mathcal{X}b)^{n-1}\mathcal{X}\right)\right] \cdot b \\
&= \left[\mu(\mathcal{X}) \cdot \mathbb{1} + \sum_{n=2}^{\infty} \widetilde{{}^c\rho_{\mu,\nu}}\left(\mathcal{X} \cdot b(\mathcal{X}b)^{n-2} \cdot \mathcal{X}\right)\right] \cdot b
\end{aligned}$$

From Theorem 4.8 and Lemma 5.8, (μ, ν) is $\boxed{c}$-infinitely divisible if and only if there exists $\mathbb{C}$-linear map $\sigma : \mathcal{B}\langle\mathcal{X}\rangle \longrightarrow \mathcal{D}$ that satisfy property (1) such that $\sigma(f(\mathcal{X})) = {}^c\rho_{\mu,\nu}(\mathcal{X}f(\mathcal{X})\mathcal{X})$ for all $f(\mathcal{X}) \in \mathcal{B}\langle\mathcal{X}\rangle$. That is

$$\begin{aligned}
{}^cR_{\mu,\nu}(b) &= \left[\mu(X) \cdot \mathbb{1} + \sum_{l=0}^{\infty} \widetilde{\sigma}(b \cdot [\mathcal{X}b]^l)\right] \cdot b \\
&= \left[\mu(X) \cdot \mathbb{1} + \widetilde{\sigma}(b[\mathbb{1} - \mathcal{X}b]^{-1})\right] \cdot b
\end{aligned}$$

hence the conclusion.

For the last part, if $(\mu, \nu) \in \Sigma^0_{\mathcal{B}:\mathcal{D}} \times \Sigma^0_{\mathcal{B}}$, then Remark 5.5 implies that $({}^c\rho_{\mu,\nu}, \rho_\nu)$ is also in $\Sigma^0_{\mathcal{B}:\mathcal{D}} \times \Sigma^0_{\mathcal{B}}$, so they satisfy the condition (2) for some $M > 0$. The representation of ${}^cR_{\mu,\nu}$ gives (for $b_1, \ldots, b_n \in M_m(\mathcal{B})$ and the identification $\mathcal{X} \cong \mathrm{Id}_m \otimes \mathcal{X}$):

$$\begin{aligned}
\|\widetilde{\sigma}(\mathcal{X}b_1\mathcal{X}b_2 \cdots b_n\mathcal{X})\| &= \|{}^c\rho_{\mu,\nu}(\mathcal{X}^2 b_1 \mathcal{X}b_2 \cdots b_n\mathcal{X}^2)\| \\
&< M^{n+3}\|b_1\| \cdots \|b_n\| \\
&< (M^2 + 1)^{n+1}\|b_1\| \cdots \|b_n\|.
\end{aligned}$$

□

References

[1] M. Anshelevich *Appell polynomials and their relatives II. Boolean theory*, Indiana Univ. Math. J. 58 (2009), 929-968.

[2] S. T. Belinschi, M. Popa, V.Vinnikov *Infinite divisibility and a non-commutative Boolean-to-free Bercovici-Pata bijection*, arXiv:1007.0058

[3] H. Bercovici, V. Pata *Stable laws and domains of attraction in free probability theory*, With an appendix by Philippe Biane, Ann. of Math. (2), **149**, (1999), no.3, 1023–1060

[4] H. Bercovici and D. Voiculescu, *Levy-Hincin type theorems for multiplicative and additive free convolution*, Pac. J. Math. 153 (1992), 217–248.

[5] F. Boca *Free products of completely positive maps and spectral sets*, J. Functional Analysis 97 (1991), 251-263

[6] M. Bozejko, M. Leinert, R. Speicher *Convolution and limit theorems for conditionally free random variables*, Pac. J. Math. 175 (1996), 357-388

[7] K. Dykema *Multilinear function series and transforms in free probability theory.* Advances in Mathematics Volume 208, Issue 1, 15 January 2007, Pages 351-407

[8] K. Dykema, E. Blanchard *Embeddings of reduced free products of operator algebras* Pacific Journal of Mathematics 199 (2001), 1-19

[9] W. Feller, *An Introduction to Probability Theory and its Applications*, Cambridge University Press, Cambridge, 1987.

[10] B. V. Gnedenko and A. N. Kolmogorov, *Limit Distributions for Sums of Independent Random Variables,* Addison-Wesley Publ. Co., Cambridge, Mass., 1954.

[11] D. S. Kaliuzhnyi-Verbovetskyi and V. Vinnikov, *Foundations of noncommutative function theory*, in preparation.

[12] A. D. Krysztek *Infinite divisibility for the conditionally free convolution*, IDAQP, Vol 10(4) (2007), 499522

[13] E. C. Lance *Hilbert C*-modules. A toolkit for operator algebraists* London Math. Soc. Lecture Note Ser. 210, 1995

[14] P. Levy, *Theeorie de L'addition des Variables Aleeatoires*, Gauthier-Villars, Paris, 1937.

[15] W. Mlotkowski *Operator-valued version of conditionally free product* Studia Mathematica 153 (1) (2002)

[16] V. Paulsen *Completely Bounded Maps and Operator Algebras.* Cambridge University Press, 2002

[17] M. Popa *Multilinear function series in conditionally free probability with amalgamation* Com. on Stochastic Anal., Vol 2, No 2 (Aug 2008)

[18] M. Popa *A new proof for the multiplicative property of the boolean cumulants with applications to the operator-valued case* Colloquium Mathematicum, Vol 117 (2009), No. 1 , 81-93

[19] M. Popa *Realization of conditionally monotone independence and monotone products of completely positive maps*, arXiv:0911.1319

[20] M. Popa *A combinatorial approach to monotonic independence over a C*-algebra*, Pac J. of Math, Vol. 237 (2008), No. 2, 299-325

[21] M. Schurmann, A. Ben-Ghorbal *Non-commutative notions of stochastic independence.* Math. Proc. Phil. Soc. (2002), 133, 531

[22] R. Speicher, A. Nica *Lectures on combinatorics of free probability*, London Math Soc Lecture Note Series 335, cambridge Univ. press 2006

[23] R. Speicher, R. Waroudi *Boolean convolution.* Fields Institute Communications, Vol. 12 (D. Voiculescu, ed.), AMS, 1997

[24] R. Speicher *Combinatorial theory of the free product with amalgamation and operator-valued free probability theory.* Memoir of the AMS 627 (1998)

[25] J.L. Taylor, *A general framework for a multi-operator functional calculus*, Advances in Math. **9** (1972), 183–252.

[26] J.L. Taylor, *Functions of several noncommuting variables*, Bull. Amer. Math. Soc. **79** (1973), 1–34.

[27] D-V. Voiculescu *Operations on certain non-commutative operator-valued random variables*, Asterisque 232 (1995), 243-275

[28] D.-V. Voiculescu *The coalgebra of the free difference quotient and free probability*, Internat. Math. Res. Not. (2000) No. 2.

[29] D-V. Voiculescu *Free analysis questions. II: The Grassmannian completion and the series expansion at the origin*, preprint, arXiv:0806.0361v1

[30] D-V. Voiculescu, K. Dykema, A. Nica *Free Random Variables* CRM Monograph Series, 1, American Mathematical Society, 1992

[31] J.-C. Wang *Limit theorems for additive c-free convolution*, Canad. J. Math. 63 (2011), no. 1, 222240

Center for Advanced Studies in Mathematics at the Ben Gurion University of Negev, P.O. B. 653, Beer Sheva 84105, Israel and
Institute of Mathematics Simion Stoilow of the Romanian Academy, P.O. Box 1-764, Bucharest, RO-70700, Romania
E-mail address: `popa@math.bgu.ac.il`

Department of Mathematics, Ben Gurion University of Negev, Beer Sheva 84105, Israel
E-mail address: `vinnikov@cs.bgu.ac.il`